

KLORA-6.3% SC

ACARICIDE AND INSECTICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Avermectin⁽¹⁾
Diamide⁽²⁾

Insecticide and Acaricide for the control of certain mites and insect pests on agricultural crops mentioned in the table.

COMPOSITION: Each liter contains:

ABAMECTIN ⁽¹⁾	1.8% W/W
CHLORANTRANILIPROLE ⁽²⁾	4.5% W/W

PROPERTIES:

KLORA- 6.3% SC: Insecticide and acaricide with a combination of (Chlorantraniliprole and Abamectin).

KLORA- 6.3% SC: Used to control mites and insects pests on agricultural crops mentioned in the table.

USES AND APPLICATION RATE :

CROP	PESTS	APPLICATION RATE ML/20 LT. WATER	PHI (DAY)	NOTES
Pome Fruit (Apple, Pear, Quince, Loquats)	Codling Moth, Light Brown Apple Moth, Native Budworm, Cotton Bollworm, Oriental Fruit Moth, Two Spotted Mite, European Red Mite	11	7	-Do not make more than one application per season. -Apply when pest reach threshold level. -Apply from 2 to 6 weeks after petal fall if monitoring shows high numbers of overwintering mites/eggs, or if mites reach threshold levels early in the season. -Do not apply to crops if a group 6 insecticide have been applied in the current season. -Do not apply to crops if three applications of a group 28 insecticide have been applied in the current season. -Do not apply KLORA-6.3% SC at rate greater than 2 X concentration
Grapes	Light Brown Apple Moth Grapevine Moth, Two Spotted Mite, Grape leaf Rust Mite, Suppression of Garden Weevil	11	28	- Apply when pest reach threshold level. - Do not make more than two application per season. - Make subsequent insecticide applications as necessary. - Do not retreat within (14) days. - Do not apply KLORA-6.3% SC at rate greater than 3 X concentration.

- For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML/20 LT. WATER	PHI (DAY)	NOTES
Cabbage	Diamondback moth	20	3	-Apply when pest reach economic threshold level and repeat as necessary.
Tomato	Spider mites, Leafminers, Caterpillars	20	3	-Days interval between applications: 5-14 days, in case of heavy pest infestation use the shorter day's interval.
Squash	Leafminers, Thrips	20	3	-Do not make more than three applications per season and then rotate with products with a different mode of action.

CAUTION:

Used only on the above-mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD: 24 hour from last application.

RE-ENTRY PERIOD OF ANIMAL GRAZING:

Do not graze or use the treated plants for stock food for 28 days after last application.

COMPATIBILITY:

Do not mix with compounds contains Kresoxim-Methyl, Fluopyram and Tebuconazole it may cause phytotoxicity to the pome fruit foliage. Compatibility test is recommended before spraying.