

ADIFON-25% WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Triazole

Systemic fungicide used to control Powdery mildew, Rusts, Black rot, Leaf blotch and Leaf spots on crops mentioned in the table of uses.

COMPOSITION: Each kg contains:

TRIADIMEFON

25% W/W (A.I)

PROPERTIES:

ADIFON-25% WP: Systemic fungicide with protective and curative action, absorbed by roots and leaves and translocated rapidly in young growing tissues.

ADIFON-25% WP: Control many fungal diseases on crops mentioned in the table of uses.

ADIFON-25% WP: Slightly toxic to mammals.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE	SAFETY PERIOD (DAYS)
Grapes	Black Rot, Powdery Mildew	150 - 200 gm/ha	14
Tomatoes, Pepper and Artichokes,	Powdery Mildew	400 - 500 gm/ha	14
Eggplant, Cucumber, Onion, Turnip	Powdery Mildew	400 gm/ha	14
Sugar Beet	Powdery Mildew, Beet Rust	500 gm/ha	14
Cereals	Powdery Mildew, Rust, Leaf Blotch, Leaf Spot	500 - 1000 gm/ha (Repeat the treatment when reinfection is observed again)	28
Peas	Powdery Mildew	200 - 300 gm/ha	14
Beans	Rust	400 - 500 gm/ha	14
Tobacco	Powdery Mildew	10 - 20 gm/100 lt. water (Repeat spray every 7 - 14 days)	14
Coffee	Rust	0.5 - 1 kg/ha (Protective spray or when first symptoms of disease appear). 1 - 2 kg/ha (In case of severe rust infection repeat every 3 - 4 weeks).	3 - 4 Weeks

Hectar= 10000 m²

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Compatible with most pesticides, spray mixture must be applied immediately after being prepared and compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.