

AZOXIFEN-32.5% SC

FUNGICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Methoxyacrylate⁽¹⁾
Triazole⁽²⁾

Fungicide with protective, curative, eradicant, translaminar and systemic properties used to control a wide range of fungal diseases on a wide range of crops such as vegetables, fruit trees, ... etc.

COMPOSITION: Each liter contains:

AZOXYSTROBIN⁽¹⁾	20.0 % W/V (A.I)
DIFENOCONAZOLE⁽²⁾	12.5 % W/V (A.I)

PROPERTIES:

AZOXIFEN-32.5% SC: Fungicide with protective, curative, eradicant, translaminar and systemic Properties used to control wide range of fungal diseases.

AZOXIFEN-32.5% SC: Slightly toxic and not toxic to environment and natural enemies.

AZOXIFEN-32.5% SC: Used in IPM control programmes.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE ML / 20 LT WATER	SAFETY PERIOD (DAYS)
Tomato and Pepper	Powdery mildew Early blight, Leaf spots	10 -15	3 7
Cucurbits except cucumber	Powdery mildew, Leaf spots and Anthracnose.	10 - 15	3
Cucumber	Powdery mildew, Downy mildew.	5	3
Potato	Early blight, Leaf holes.	10 - 15	14
Peach	Powdery mildew, Leaf spots, Leaf holes.	10 - 15	14
Grapes	Powdery mildew, Leaf spots. Downy mildew, Dieback	10 - 15 10	14

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Some types of apple are sensitive to **AZOXIFEN-32.5% SC** so avoid spray drift to neighboring apple orchard it may cause severe burns.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Compatible with most pesticides, use the solution directly after mixing, and compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.