

BROMOSTAR-50% EC

ACARICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Diphenylcarbinol

Non-systemic acaricide with contact action, and long residual activity.

COMPOSITION: Each Liter contains:

BROMOPROPYLATE

50% W/V (A.I)

PROPERTIES:

BROMOSTAR-50% EC: Effective to control of all stages of Tetranychid and Eriophyid mites on the agricultural crop listed in the table of uses.

BROMOSTAR-50% EC: Inhibitor of oxidative phosphorylation, disruptor of ATP formation (inhibitor of ATP synthase).

USES AND APPLICATION RATE:

CROP	PEST	APPLICATION RATE / 20 L OF WATER	SAFETY PERIOD (DAYS)
Apples, pears, plum, apricot, cherry, almond, peach, nectarine	Bud mite, eriophyid mites	10 – 15 ml	21
	Red spider mite, flat mite	15 - 20 ml	
Oranges, lemon, grapefruit, mandarin.	Bud mite, eriophyid mites	10 – 15 ml	21
	Red spider mite, flat mite	15 - 20 ml	
Tomato, potato, eggplant, pepper, cucumber, squash	Red spider mite	20 - 30 ml	14
Ornamentals	Red spider mite	15 - 20 ml	---

CAUTION:

Used only on the above mentioned crops, and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours from last application.

COMPATIBILITY:

The product is compatible with most products except alkaline materials such as copper, sulfur or lime sulfur compounds, compatibility test is recommended before spraying

• For more details about first aid & precautions please refer to first aid & precautions index.