

CRIPTAN-50% WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Phthalimide

Fungicide with protective and curative action controls a wide range of fungal diseases on a wide range of crops.

COMPOSITION: Each kg contains:

CAPTAN

50% W/W (A.I)

PROPERTIES:

CRIPTAN-50% WP: Used to control various fungal diseases affecting vegetation and seedlings.

CRIPTAN-50% WP: Control many of fungal diseases, affecting crop plants such as spots, leaf curl, late and early blight and seedlings diseases.

CRIPTAN-50% WP: With low mammalian toxicity.

USES AND APPLICATION RATE:

Apply by spraying of the plant or on green parts as follow:

CROP	DISEASE	APPLICATION RATE GM /20 LT. WATER	APPLICATION METHOD	APPLICATION RATE PER SEASON	SAFETY PERIOD (DAYS)
Apple, Pears	Scab, Rot, Blotch, Spots	20-40	Spraying	Apple: 726 gm./ 20 Lt. Water	7
Apricot, Plum, Peach, Cherry.	Brown Mold, Leaf Spot, Scab	40-50	Spraying	Apricot: 284 gm 20 Lt. Water Plum: 612 gm/20 Lt. Water Peach: 726 gm 20 Lt. Water Cherry: 318 gm 20 Lt. Water	3 except Cherry 5
Grapes	Dieback. Black Mold, Downy Mildew.	40 40-50	Spraying Spray before, after flowering and at the beginning of setup. Apply after 2-3 weeks when downy mildew occur.	272 gm/20 Lt. Water	7
Strawberry	Grey Mold, Leaf spot	90-135	Spray in the beginning of growth in spring and repeat the spraying every week	544 gm/20 Lt. Water	3

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE GM. /20 LT. WATER	APPLICATION METHOD	APPLICATION RATE PER SEASON	SAFETY PERIOD (DAYS)
Cucumber, squash, pumpkin	Anthracnose, Scab	65-135	Spray at the beginning of disease and repeat every 5-7 days	-----	3
Tomato	Anthracnose, Early & Late Blight, Septoria Leaf Spot	40-80	Spray at the beginning of disease and repeat every 7-10 days	-----	3
Broccoli, Cabbage, Cauliflower, Eggplant, Pepper, Tomato, Rose, Peas	Seedling Wilt, Root Rot	125-225 gm. /100 m ²	Applied to soil before planting at 8-10 cm	-----	-----
Ornamentals	Leaf Spot, Blight, Black Spot	40-45	Spraying Repeat every 7-10 days if needed	-----	-----

CAUTION:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

NOTE:

Can be used for peas seed treatment to control soil disease as 160 gm **CRIPTAN-50% WP** / 500 ml water/100 kg seed.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Incompatible with all alkaline compounds, Oil sprays, TEPP compound, Emulsifiable Parathion compounds.

• For more details about first aid & precautions please refer to first aid & precautions index.