

CURTINE-V WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Dithiocarbamate⁽¹⁾
cyanoacetamide oxime⁽²⁾

Effective fungicide to control a broad spectrum of fungal diseases such as downy mildew and blights.

COMPOSITION: Each kg contains:

| | |
|--------------------------|----------------|
| MANCOZEB ⁽¹⁾ | 46.5%W/W (A.I) |
| CYMOXANIL ⁽²⁾ | 4 %W/W (A.I) |

PROPERTIES:

CURTINE-V WP: Highly absorbed by plants, ensuring maximum utilization of the pesticide before begin glossed by rain.

CURTINE-V WP: Protective fungicide, prevent the formation of Mycelium at the incubation period for 4 days after infection.

CURTINE-V WP: Could be sprayed three days after rain to maintain good results.

USES AND APPLICATION RATE:

| CROP | DISEASE | APPLICATION RATE / 20 LT. WATER | SAFETY PERIOD (DAYS) |
|---------------|-----------------------|------------------------------------|-------------------------|
| Grapes | Downy mildew | 40-50 gm | 14 |
| Potatoes | Blights, Downy mildew | 40-50 gm | 15 |
| Onion, Garlic | Downy mildew | 40-50 gm | 14 |
| Tobacco | Downy mildew | 40-50 gm | 14 |

CAUTION:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hour from last application.

COMPATIBILITY:

Compatible with most the pesticides, but compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.