

CYMODON- 52.5% WDG

FUNGICIDE
WATER DISPERSIBLE GRANULES

VAPCO

Class:
Cyanoacetamide Oxime⁽¹⁾
Oxazolidinedione⁽²⁾

Is a systemic, contact and antisporent fungicide for controlling fungal diseases on crops listed in the table of uses.

COMPOSITION: Each Kg contains:

CYMOXANIL ⁽¹⁾	30.0% W/W (A.I)
FAMOXADONE ⁽²⁾	22.5% W/W (A.I)

PROPERTIES:

CYMODON- 52.5%WDG: is a systemic and contact fungicide used to control many fungal diseases on the crops listed on the table of uses.

CYMODON- 52.5%WDG: it is composed of the association of two active materials, famoxadone and cymoxanil.

CYMODON- 52.5%WDG: famoxadone inhibits mitochondrial electron transport and spore germination while cymoxanil a penetrant anti-mildew with a curative action.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE / 20 LITER WATER	MAX. NUMBER OF TREATMENTS	APPLICATION TIME	SAFETY PERIOD (DAYS)
Potato	Late Blight, Early Blight	8 gm	4	Pre-blooming up till harvest	14
Tomato	Downy Mildew, Alternaria				13
Lettuce					
Cucumber, Squash, Watermelon, Muskmelon Pumpkin					

CAUTION:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Not compatible with alkaline material and compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.