

FOLDAN-50% WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Phthalimide

A protective fungicide with contact action, effective in controlling many of the most fungal diseases on a wide range of crops.

COMPOSITION: Each kg contains:

FOLPET

50% W/W (A.I)

PROPERTIES:

FOLDAN-50% WP: A contact fungicide with a curative and protective action.

FOLDAN-50% WP: Effective to control many of the most economically devastating fungal diseases on crops mentioned in the tabel of uses.

FOLDAN-50% WP: Slightly toxic to mammals and other beneficial insects or wildlife.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE / 20 LT. WATER	NUMBER OF APPLICATION PER SEASON	SAFETY PERIOD (DAYS)
Pear	Scab, Leaf Spot, Powdery Mildew	50 gm	10	21
Apple		40 gm	Every 14 days	21
Citrus	Anthracoese, Dieback, Gummosis	60 gm	---	21
Grapes	Downy Mildew, Powdery Mildew, Brown Rot, Dieback.	60 gm	Every 14 days	8
Apricot, Plum, Peach, Almond	Leaf Curl, Anthracnose, Powdery Mildew, Leaf Spot.	50 gm	3	10
Olives	Peacock Eye Spot, Wilt	30 gm	---	10
Cucumber, Squash, Pumpkin	Powdery Mildew, Downy Mildew, Stem Rot, Anthracnose.	15 gm	---	7
Eggplant, Peas	Powdery Mildew, Early Blight, Late Blight	30 gm	---	10
Legumes	Powdery Mildew, Leaf Spot	7 - 8 gm	---	10
Lettuce	Downy Mildew, Gray Mold	40 gm	Weekly	11

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE / 20 LT. WATER	NUMBER OF APPLICATION PER SEASON	SAFETY PERIOD (DAYS)
Muskmelon	Downy Mildew, Powdery Mildew, Stem Rot, Anthracnose.	40 gm	Weekly	7
Watermelon	Downy Mildew, Powdery Mildew, Stem Rot, Anthracnose.	50 gm	1 - 2	7
Onions	Purple Blotch, Neck And Leaf Rot.	40 - 50 gm	2 - 3	7
Pepper	Wilt, Stem Rot, Powdery Mildew.	10 - 30 gm	3 - 4	7
Potatoes	Early Blight, Late Blight	60 gm	Weekly	14
Tomatoes	Early Blight, Late Blight	40 gm	Weekly	14

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

PHYTOTOXICITY:

Not phytotoxic, except for: Sweet Cherries, D'Anjou variety of Pears and sensitive Apple varieties is possible if applied early in cropping before harvest.

RE-ENTRY PERIOD:

24 hours from the last application.

COMPATIBILITY:

Compatible with most pesticides except highly alkaline compounds. It's preferable to use the product immediately after mixing and compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.