

HAWK-25% EC

HERBICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Hydroxybenzonitrile

A selective contact herbicide, used as crops post-emergence to control a wide range of annual broadleaf weeds as shown in the table of uses.

COMPOSITION: Each Liter contains:

IOXYNIL OCTANOATE

25% W/V (A.I)

PROPERTIES:

HAWK-25% EC: Is a contact herbicide which penetrates through the leaves and translocated to the root and inhibits photosynthesis.

HAWK-25% EC: A selective herbicide with some systemic activity to control annual broadleaf weeds.

USES AND APPLICATION RATE :

CROP	RATE OF APPLICATION	TIME OF APPLICATION
Onion, garlic and leek.	200-280 ml/donum	Form 3 leaf stage.
Wheat, barley.	160-200 ml/donum	Form 3 leaf stage to stem elongation, it's preferable during tillering stage.

CAUTION:

Used only on the above mentioned crops.

Donum = 1000m²

REMARKS:

- Calibration of sprayer is important before application.
- Use high dosage rate for spraying when weed infestation is heavy.
- Do not use spray equipment contaminated with this product for other purposes unless thoroughly cleaned with water and detergent solution.
- In order to achieve maximum control of the targeted weeds, apply **HAWK-25% EC** to actively growing weeds. The weeds are best controlled at emergence and at 3 true leave stage.

RE-ENTRY PERIOD:

48 hours from the last application.

COMPATIBILITY:

Compatible with most herbicides, but incompatible with liquid fertilizers and water soluble formulations of 2,4-D and it's preferable to make compatibility test before spraying.

* **OTHER TRADE NAME: HOCKS-25% EC.**

• For more details about first aid & precautions please refer to first aid & precautions index.