

HERBSHOCK-24% EC

HERBICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Aryloxyphenoxypropionate

A selective systemic herbicide used to control wild oats and canary grass in wheat fields.

COMPOSITION: Each litre contains:

CLODINAFOP PROPARGYL 24% W/V (A.I)

PROPERTIES:

HERBSHOCK-24% EC: Selective systemic herbicide should be applied only when weeds are actively growing as the controlling symptoms appears from 1-3 weeks after application and it affect the meristematic tissue.

USES AND APPLICATION RATE :

CROP	WEED	APPLICATION RATE	TIME OF APPLICATION		SAFETY PERIOD (DAYS)
			CROP	WEEDS	
Wheat	Wild oats	230 ml / ha	Before the appearing the fourth leaf at the tillering stage	At 1 to 6 leaf stage on the main stem when weed growth is active	60
	Canary Grass			At 1 to 5 leaf stage on the main stem when weed growth is active	

Hectare = 10000 m²

Donum = 1000 m²

CAUTION:

Used only on the above mentioned crop and follow the recommended safety period before harvest.

REMARKS:

- The pesticide is diluted with an amount of water not less than 100-200 liters/ha.
- Do not spray until two hours after the rain
- Do not spray when the plant is in a state of stress, such as when the soil is dry, high humidity, lack of elements, insect or disease infestations, or when frost occurs.
- The sprayer must be calibrated before application.
- The spray must be even and homogeneous throughout the area to be treated.
- It is not recommended to harvest the crop before 60 days of treatment.
- Avoid applying the pesticide to barley or any other crop other than wheat. The spray is not allowed to drift or fly into neighboring fields planted with crops other than wheat
- The pesticide can be mixed with a non-ionic adjuvant as per the ratios shown in the adjuvant usage table.

• For more details about first aid & precautions please refer to first aid & precautions index.

RE-ENTRY PERIOD:

24 hours from the last application.

GRAZING RESTRICTIONS

Do not graze or feed animals for 72 hours from the last application.

COMPATIBILITY

* It can be mixed with other pesticides as long as all mixing instructions mentioned in the labels of the pesticides to be mixed with **HERBSHOCK-24% EC** are followed. And follow the spraying precautions, restrictions, crop rotation, application timing and pre-harvest periods mentioned in the labels of the pesticides that will be mixed with **HERBSHOCK-24% EC**.

* An initial compatibility test must be done before mixing with other herbicides on a small scale to avoid any sensitivity of the crop.

• For more details about first aid & precautions please refer to first aid & precautions index.