

# NOGREEN-28% EC

HERBICIDE  
EMULSIFIABLE CONCENTRATE

VAPCO

Class:  
Hydroxybenzotrile

Selective contact herbicide with some systemic activity, to control post-emergence annual broad-leaved weeds in wheat, barley, maize, onions, garlic, alfalfa, non-crop land and industrial sites.

**COMPOSITION:** Each Liter contains:

**BROMOXYNIL OCTANOATE**

**28% W/V (A.I)**

**PROPERTIES:**

**NOGREEN-28% EC:** Selective herbicide with some systemic activity, absorbed by the foliage, with limited translocation.

**NOGREEN-28% EC:** Controls weeds by inhibits photosynthesis.

**NOGREEN-28% EC:** Selective herbicide used to control post-emergence annual broadleaf weeds in wheat, barley, maize, onions, garlic, alfalfa, non-crop land and industrial sites.

**NOGREEN-28% EC:** Moderately toxic to mammals, toxic to fish and not toxic to bees.

**USES AND APPLICATION RATE :**

CROPS	WEEDS	DOSAGE AND APPLICATION TIME	SAFETY PERIOD
Wheat, Barley	Annual broadleaf weeds such as: Chrysanthemum, Galium, Prickly dock.	1.5 - 2 lt./ha (200 - 400 lt. of water/ha) Post-emergence to the beginning of tillering stage.	42 days
Maize		1.5 - 2 lt./ha (200 - 400 lt. of water/ha) Apply at 2 - 8 leaf stage.	42 days
Garlic		2.3 - 3 lt./ha (200 - 400 lt. of water/ha) Apply post-emergence but before 30 cm height.	3.5 months
Onions		1.5 - 2 lt./ha (200 - 400 lt. of water/ha) Apply at 2 - 5 leaf stage.	1 month
Alfalfa		1.5 - 2 lt./ha (200 - 400 lt. of water/ha) Apply when weeds at 5 cm height.	1 month
Non-crop land and industrial sites		1.5 - 2 lt./ha (200 - 400 lt. of water/ha) Apply when weeds have emerged and are actively growing.	---

Donum = 1000 m<sup>2</sup> Hectare = 10000 m<sup>2</sup>

**CAUTION:**

Used only on the above mentioned crops and follow the recommended safety period before harvest.

• For more details about first aid & precautions please refer to first aid & precautions index.

**REMARKS:**

- Calibration of sprayer is important before application.
- Weeds should be sprayed before 6 leaf stage.
- The Spray should be homogeneous in all treated areas.

**GRAZING RESTRICTIONS:**

Do not graze or feed animals at least 60 days from the last applications.

**RE-ENTRY PERIOD:**

48 hours from the last application.

**COMPATIBILITY:**

Compatible with most herbicides and it's preferable to make compatibility test before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.