

# OSCAR-75% WP

HERBICIDE  
WETTABLE POWDER

VAPCO

Class:  
Sulfonylurea

Post-emergence herbicide used to control broadleaf weeds on Wheat, Barley, Oats and Maize.

**COMPOSITION:** Each Kg contains:

TRIBENURON – METHYL

75% W/W (A.I)

## PROPERTIES:

**OSCAR-75% WP:** A foliar acting sulfonylurea herbicide.

**OSCAR-75% WP:** Rapidly absorbed by foliage and roots and translocated throughout the plant.

**OSCAR-75% WP:** Post-emergence herbicide, control broadleaf weeds on wheat and barley, the effect of the pesticide begins almost immediately after treatment and symptoms can be seen in about 2 weeks after spraying.

## USES AND APPLICATION RATE :

CROP	WEEDS	TIME OF APPLICATION	APPLICATION RATE
Wheat	Pimpernel, Chickweed, Shepherd`s Purse, Spurrey, Californian Thistle, Wild Turnip, Willow Weeds, Scotch Thistle and Fathen.	Spray after the crops in two leaf stage, but before the boot stage.  Incuse of severe weed injury and large weeds size and density.	17 gm / ha
Barley	Pimpernel, Chickweed, Shepherd`s Purse, Spurrey, Californian Thistle, Wild Turnip, Willow Weeds,		25 gm / ha
Oats	Amaranthus, Sheep`s Sorrel, Twin Cress, Wire Weeds, Scotch Thistle, Fathen and Redroot.		35 gm / ha
Maize	Broadleaf weeds	When crops are 20-30 cm high	1.0 – 1.5 gm / donum

Donum = 1000m<sup>2</sup>

Hectare = 10000m<sup>2</sup>

**CAUTION:** Used only on the above mentioned crops.

**SAFETY PERIOD:** All crops can be planted 60 days after treatment, but wheat can be planted any time after treatment.

## APPLICATION METHOD:

- To obtain best result use ground sprayer but be sure that the amount of water not less than (100 – 300 lt./ha).
- In Maize don't use more than 1.5 gm/donum to avoid severe crop injury.

• For more details about first aid & precautions please refer to first aid & precautions index.

**NOTE:**

Adding spreading and sticking agent like ( **SUPER-SERFIX** ) can increase **OSCAR-75% WP** efficiency.

**REMARKS:**

- Calibration of sprayer is important before application.
- Complete spray coverage must be achieved.
- Maximum number of treatment 1 per season.
- Special care must be taken to avoid damage by drift into near crops or water sources.
- Do not apply to wheat or barley crops planted with another crops.
- All crops can be planted 60 days after treatment, but Wheat can be planted immediately after treatment.
- Do not apply under unfavorable conditions like drought, low fertility, water saturated soil, disease or insect damage, and under certain conditions such as prolonged cool weather, wide fluctuation in day / night temperatures just prior to or soon after treatment, temporary yellowing and / or stunting may occur.
- Addition of surfactants (e.g. **SUPER SURFIX**) significantly improves the efficacy of **OSCAR-75% WP**.
- Increasing levels of soil moisture and warmer temperatures increase the pesticide efficacy.
- The spray solution must be used during 24 hours with continuous agitation to avoid degradation and keep efficiency.

**GRAZING RESTRICTIONS:**

Do not graze or feed animal from treated area for 60 days from the last application.

**RE-ENTRY PERIOD:**

24 hours from the last application.

**COMPATIBILITY:**

- Compatible with most pesticides, but it is preferable to make compatibility test before spraying.
- **OSCAR-75% WP** mixed with **VAPCOR-70% WP** (Metribuzin) may reduce the control of wild Garlic.
- **OSCAR-75% WP** mixed with (Dicamba) containing compounds may reduce the control of some broadleaf weeds.
- Don't mix **OSCAR-75% WP** with (Diclofop-methyl) as grass control may be reduced.
- Don't mix **OSCAR-75% WP** with Organophosphorus insecticide (such as Malathion) as crop injury will result. Temporary crop yellowing or in severe cases crop injury.
- Do not use **OSCAR-75% WP** in liquid fertilizer solution having pH of less than 3, conduct a small compatibility test before spraying.

**TRADE NAME: OSCARD – 75% WP**

- For more details about first aid & precautions please refer to first aid & precautions index.