

P-OXYRIDE 50% WP

FUNGICIDE
WETTABLE POWDER

Class:
Copper Compound

A protective fungicide with contact action. Effective in controlling many of fungal diseases on a wide range of agricultural crops .

COMPOSITION : Each kg contains:

COPPER OXYCHLORIDE 50% W/W (A.I)

PROPERTIES:

P-OXYRIDE 50% WP: A protective foliar fungicide with contact action, deposits must be on the crop before fungal spores begin to germinate.

P-OXYRIDE 50% WP: Control a wide range of fungal diseases on a wide range of Crops.

P-OXYRIDE 50% WP: Not phytotoxic for treated plants , but may cause injury to plum and peach foliage, especially at low temperatures.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE / 20 LT. WATER	APPLICATION TIME AND METHOD
Apricot Plum, Cherry, Peaches, Almond.	Shot hole, Leaf Curl.	50 gm	- Apply at bud swell and with in one week of bud opening . - Repeat 7 - 10 days .
Apple, Pears	Scab	50 gm	- Repeat 10 - 14 days.
Grapes	Downy Mildew	50 gm	- First apply when shoots are 10 cm long, repeat at 10 - 14 day.
Citrus	Black Spot, Melanose, Smoky Blotch, Scab On Lemon.	40-60 gm	- Add 30 ml of summer oil. (apply at 50 - 75% petal fall. use higher rate in coastal districts)
Potatoes	Early Blight, Late Blight	45-50 gm	- Apply from crop emergence to crop maturity at 7 - 14 day .
Tomatoes	Early Blight, Late Blight, Bacterial Spot, Septoria Spot Canckers	45-50 gm	- Seedbeds: apply each 7 days under suitable humidin conditions and repeat at 7 - 14 day .

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE / 20 LT. WATER	APPLICATION TIME
Cucumber, Watermelon, Cucurbit, Squash	Bacterial Spot, Angular Leaf Spot	50 gm	- Repeat 10 - 14 days.
Cabbage, Cauliflower	Bacterial Spot, Downy mildew, Black Leaf Spot	50 gm	- This product must not be applied to cabbage, when frost is likely to occur. - Repeat 10 - 14 days .
Lettuce	Bacterial Spot, Downy mildew, Anthracnose.	50 gm	- This product must not be applied to lettuce, when frost is likely to occur. - Repeat 7 - 10 days .
Carrot	Leaf Spot	50 gm	- Repeat 7 - 14 days .
Celery	Early Blight, Leaf Spot	50 gm	- Repeat 7 - 14 days .
Onion	Downy Mildew	50 gm	- Repeat 10 days .
Beans	Rust, chocolate spot	50 gm	- Repeat 10 - 14 days .
Peas	Blight	50 gm	- Repeat 10 - 14 days .
Ornamentals	Fungal Leaf Spots, Downy mildew	60 gm	- Repeat 7 - 14 days. - Small scale phytotoxicity test are recommended as some varieties may be sensitive.

SAFETY PERIOD:

1 day for the above mentioned crops.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Compatible with most pesticides except alkaline materials like lime sulfur and compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.