

PENDILIN- 40% SC

HERBICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Dinitroaniline

Selective herbicide used to control annual grasses and broadleaf weeds on a wide range of crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

PENDIMETHALIN

40 % W/V (A.I)

PROPERTIES:

PENDILIN- 40% SC: Selective herbicide absorbed by leaves and roots.

PENDILIN- 40% SC: Used against most annual grasses and broadleaf weeds.

PENDILIN- 40% SC: Applied as pre-plant, pre-emergence, pre-transplanting, or early post-emergence on a wide range of crops which are mentioned in the table of uses.

USES AND APPLICATION RATE :

CROPS	TIME OF APPLICATION	APPLICATION RATE (LT./HA)	COMMENTS
Winter barley, winter wheat	Pre-crop emergence	3.3	Do not apply to crops after 30th November.
Spring barley	Pre-crop emergence	3.3	- Apply after planting and before emergence. - Do not apply PENDILIN- 40% SC after the end of March unless rainfall is imminent.
Onion, leek	Pre-crop emergence Pre-transplanting	3.3	- Apply after planting but before crop or weed emergence. - Transplanted crops (apply pre-transplanting) - Do not apply to any onion or leek crop when heavy rain is expected.
Carrot	Pre-crop emergence	3.3	- Apply after planting but before crop and weed emergence.
Broccoli, cabbage, cauliflower	Before transplanting	3.3	- Apply after final plant bed cultivation but before transplanting - Do not apply PENDILIN- 40% SC post-planting as crop damage may occur. - Do not apply PENDILIN- 40% SC to any transplanted brassicas when heavy rain is expected.

• For more details about first aid & precautions please refer to first aid & precautions index.

CROPS	TIME OF APPLICATION	APPLICATION RATE (LT./HA)	COMMENTS
Potato	Pre-crop emergence	3.3	- Apply immediately after planting. - Do not apply later than 7 days before emergence.
Strawberry	After flower initiation but before flower trusses emergence	3.3	- PENDILIN- 40% SC should be applied to dormant newly planted in the autumn or early spring immediately after planting and prior to weed emergence. - Pre-planting application of PENDILIN- 40% SC is not recommended.
Apple, cherry, pear, plum	Before bud burst	3.3	- Apply PENDILIN- 40% SC from autumn to early spring. - PENDILIN- 40% SC is not recommended for use after the end of March unless adequate soil moisture is present

Hectare = 10000m²

CAUTION:

Used only on the above mentioned crops.

REMARKS:

- Do not apply **PENDILIN- 40% SC** to crops suffering from stress.
- Seed should be covered with a minimum of 2 - 3 cm of settled soil.
- Do not disturb the soil after **PENDILIN- 40% SC** has been applied as this will reduce weed control.
- Do not spray if the seeds are scattered.
- **PENDILIN- 40% SC** should not be used on protected crops or in greenhouses.
- Do not use on soils with more than 10% organic matter.
- Some soil moisture must be present for **PENDILIN- 40% SC** to be activated. Best results will be obtained if rainfall occurs within seven days of application.
- All dosages should be applied with 100-200 liters of water per hectare.

RE-ENTRY PERIOD:

24 hours from the last application.

GRAZING RESTRICTIONS:

Do not graze or feed animals from treated area for 24 hours from the last application.

COMPATIBILITY:

Incompatible with other pesticides and fertilizers.

- For more details about first aid & precautions please refer to first aid & precautions index.