

# PROPORID-72.2% SL

FUNGICIDE  
SOLUBLE CONCENTRATE

VAPCO

Class:  
Carbamate

Systemic fungicide with preventive and curative action used to control soil born fungi and fungal diseases on crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

PROPAMOCARB HYDROCHLORIDE

72.2% W/V ( A.I)

#### PROPERTIES:

**PROPORID-72.2% SL:** Systemic fungicide with preventive and curative protective action used to control soil born fungi and fungal diseases on crops mentioned in the table of uses.

**PROPORID-72.2% SL:** Has long residual activity for 3 - 8 weeks depends on temperature, lights and soil type.

**PROPORID-72.2% SL:** Low toxic to bees and fish.

#### USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE	SAFETY PERIOD (DAYS)
Tomato, Cabbage, Raddish and Potato	Root rot ( <i>Pythium</i> and <i>Phytophthora</i> )  Downy mildew  Late blight ( <i>Phytophthora</i> )	<b>Seed treatment:</b> Directly after planting, add 70 ml/20 Lt. water to the soil surface at the rate of 2 - 4 Lt. (solution)/m <sup>2</sup> . <b>Nursery treatment::</b> • <b>Before transplanting:</b> add 70 ml/20 Lt. water to the soil surface at the rate of 2 - 4 Lt. (Solution)/m <sup>2</sup> . • <b>After transplanting:</b> add 20ml / 20 Lt. water to the soil surface at the rate of 200 ml (Solution) / plant, repeating every two weeks as necessary. <b>Foliar spray:</b> 30 ml / 20 Lt. water	7 except cabbage 14
Cucumber	Root rot ( <i>Pythium</i> and <i>Phytophthora</i> )  Downy mildew	<b>Seed treatment:</b> Directly after planting, add 70 ml /20 Lt. water to the soil surface at the rate of 2 - 4 Lt. (Solution)/m <sup>2</sup> . <b>Nursery treatment:</b> • <b>Before transplanting:</b> Add 70 ml /20 Lt. water to the soil surface at the rate of 2 - 4 Lt. (Solution)/m <sup>2</sup> . • <b>After transplanting:</b> Add 20 ml / 20 Lt. water to the soil surface at the rate of 200 ml (Solution) / plant, repeating every two weeks as necessary. <b>Foliar spray:</b> 50 ml / 20 Lt. water	3

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE	SAFETY PERIOD (DAYS)
Ornamental	Root rot ( <i>Pythium and Phytophthora</i> )	<b>Seed Planting and Nursery:</b> Uses and application rate same as mentioned before. <b>Cutting:</b> Add 5 - 15ml /10 Lt. water to the soil surface at the rate of 5 - 10 Lt. (Solution) /M <sup>2</sup> .	---

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

24 hours from last application.

**COMPATIBILITY:**

Compatible with most common used pesticides, incompatible with liquid fertilizers, it's recommended use directly after mixing and compatibility test recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.