

RIMANIL-40% SC

FUNGICIDE
SUSPO-EMULSION

VAPCO

Class:
Anilinopyrimidine

Systemic fungicides penetrate the leaves cuticle and reach the plant tissue to inhibit fungi and spores germination.

COMPOSITION: Each liter contains:

PYRIMETHANIL

40% W/V (A.I)

PROPERTIES:

RIMANIL-40% SC: systemic fungicides penetrate the leaves cuticle and reach the plant tissue to inhibit fungi and spores germination.

RIMANIL-40% SC: can be used in tomato greenhouses.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE / 20 LITER WATER	NOTES	SAFETY PERIOD (DAYS)
Grape	Gray Mold	40 ml 21 ml When mixing with other pesticides	- Use sufficient water to ensure penetration and coverage for flowers and bunches - Spray at early bloom stage, version stage and pre-harvest.	14
Apricot Nectarine Peach, Plum (do not apply on cherries)	Brown Rot, Blossom Blight	21- 40 ml	- (Do not apply on cherries) - Begin application when condition becomes favorable for disease. - Do not apply more than three applications, days interval between application not less than 7 days	14
	Shot Hole, Scab, Gray Mold	40 ml 21 ml When mixing with other pesticides	- Apply at full bloom stage or at petal fall stage - Pre-harvest application should be made.	
Tomato	Early Blight Gray Mold	15 ml	- Begin application when condition becomes favorable for disease. - Days interval between application 7-14 days - Greenhouse use: apply in well-ventilated plastic houses or glass houses. Ventilate for at least 2 hours after application.	7
Potato, Sweet Potato	Early Blight	15 ml	- Begin application when condition becomes favorable for disease. - Days interval between application 7-14 days	7

• For more details about first aid & precautions please refer to first aid & precautions index.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE/20 LITER WATER	NOTES	SAFETY PERIOD (DAYS)
Onion Garlic Leek	Leaf Blight, Neck Rot, Purple Blotch	40 ml 21 ml When mixing with other pesticides	- Begin application when condition become favorable for disease. - Days interval between application 7-14 days	7
Strawberry	Gray Mold	40 ml 21 ml When mixing with other pesticides	- Make applications from pre-bloom stage to harvest on a 7 -14 days interval depending upon disease conditions.	7

CAUTION:

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Do not apply on cherries.
- Add half of the required water to the tank then start agitation, during agitation add **RIMANIL -40% SC** and then add the rest required water (maintain agitation during application) .

CROP TOLERANCE:

May be phytotoxic in closed systems at $\geq 80\%$ on certain species, ventilation is recommended to avoid leaf spots on certain crops.

RE-ENTRY PERIOD:

24 hours from last application for grapes and 12 hours from last application for other crops.

COMPATIBILITY:

Compatible with most herbicide, insecticide, and foliar nutrient products, fungicides (except **CAPTAN-50% WP**), and compatibility test is recommended before application.

- For more details about first aid & precautions please refer to first aid & precautions index.