

# SPIDERON-10% WP

ACARICIDE  
WETTABLE POWDER

VAPCO

Class:  
Carboxamide

Non systemic acaricide with contact action, and long residual activity, used to control many species of mite on target crops.

COMPOSITION: Each Kg contains:

HEXYTHIAZOX

10% W/W (A.I)

## PROPERTIES:

**SPIDERON-10% WP:** Non systemic acaricide with contact action.

**SPIDERON-10% WP:** Highly effective in controlling mites in eggs and nymph stage that mean complete controlling for mites.

**SPIDERON-10% WP:** has long residual activity, non-harmful to beneficial insects.

## USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATES / 20 LT. WATER	SAFETY PERIOD (DAYS)
Lemon, Orange, pomelo, mandarin, grapefruit	European Red Mite.	8 - 10 gm	7
Apple	Red Spider Mite, Two-spotted Spider Mite.	8 - 10 gm	7
Peaches, Grapes, Pear, Apricots	Red Spider Mite	8 - 10 gm	7
Strawberries	Red Spider Mite	8 - 10 gm	3
Fig, Eggplant, Peppers, Peas, Cucumber	Two-spotted Spider Mite.	8 - 10 gm	3
Beans	Red Spider Mite	8 - 10 gm	14
Ornamentals	Red Spider Mite	8 - 10 gm	-----

## Caution:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

## RE-ENTRY PERIOD:

24 hours from last applications.

## COMPATIBILITY:

Compatible with most common pesticides, including alkaline compounds, compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.