

# TAKAMOL-4% EC

FUNGICIDE  
EMULSIFIABLE CONCENTRATE

VAPCO

Class:  
Pyrimidine Carbinol

Systemic fungicide with protective, curative and eradicant action.

**COMPOSITION:** Each liter contains:

FENARIMOL

4 % W/V (A.I)

**PROPERTIES:**

**TAKAMOL-4% EC :** Systemic fungicide with protective, curative and eradicant action to control powdery mildew on different crops.

**TAKAMOL-4% EC :** Stable, not affected by rain and high temperatures.

**TAKAMOL-4% EC :** Does not affect pollination and setting of plants.

**USES AND APPLICATION RATE:**

CROP	DISEASE	APPLICATION RATE /20 LT WATER	NOTES	SAFETY PERIOD (DAYS)
Apples	Powdery Mildew.	15 ml	The amount of water per donum 150 - 200 lt.	14
	Scab	18 ml	Repeat spray every 14 days.	14
Cucumber, Watermelon, Muskmelon, Squash, Pumpkins.	Powdery Mildew.	5 ml	The amount of water per donum 100 lt. Repeat spray every 14 days.	4

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**REMARKS:**

- Do not use **TAKAMOL-4% EC** through any irrigation system.
- The product will not affect the buds, leaves & fruits if used as recommended but if used at high dose that will lead to stunt, abnormal leaves growth & increase in coloring of leaves.

**RE-ENTRY PERIOD:**

24 hours from last application.

**COMPATIBILITY:**

Compatible with most pesticides, but compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.