

TEBIZOLE-25% WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Triazole

Systemic fungicide with protective and curative action to control many fungal diseases on a wide range of crops and to protect seed during germination stage.

COMPOSITION: Each kg contains:

TEBUCONAZOLE

25 % W/W (A.I)

PROPERTIES:

TEBIZOLE-25%: Systemic fungicide with protective and curative action.

TEBIZOLE-25%: Fungicide recommended to control diseases such as Rust, Powdery mildew, Early leaf spot, Late leaf spot, Net blotch, Stem and pod rot, Rhizoctonia wilt, Monilia, Scaled, Scab, Septoria spot, Early blight, Stem canker on Almond, Eggplant, Barley, Bean, Cucumber, Grapes, Peanut, Plum, Potatoes, Sunflower, Sweet pepper, Tomatoes, Wheat, Cereal and Corn.

TEBIZOLE-25%: Effective against various Smut and Bunt diseases of Cereals.

TEBIZOLE-25%: Slightly toxic to Bees

USES AND APPLICATION RATE:

A.FOLIAR APPLICATION:

CROP	DISEASE	APPLICATION RATES GM / HA	SITE	NO. OF APPLICATIONS	SAFETY PERIOD (DAYS)
Almond	Monilia, powdery mildew.	200 gm/ha (20 gm/200 lt. Water)	Field	1	21
Eggplant	Powdery mildew.	600 gm/ha (60 gm/200 lt water)	Field & green house	3	14
Barley	Rust, powdery mildew, scald, head scab, septoria spot.	752 gm/ha (75 gm/200 lt water)	Field	1	30
Bean	Leaf spot, rust	1.0 kg/ ha (100 gm/200 lt water)	Field	1-2	14
Grapes	Powdery mildew	600 gm/ha (60 gm/200 lt water)	Field	2	14
Walnut	Early leaf spot, late leaf spot, rust, net blotch.	1.0 kg/ ha (100 gm/200 lt water)	Field	1	21

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATES GM / HA	SITE	NO. OF APPLICATIONS	SAFETY PERIOD (DAYS)
Plum	Monilia, powdery mildew.	200 gm/ha (20 gm/200 lt water)	Field	1	21
Sunflower	Stem canker, stem rot.	500 gm/ha (50 gm/200 lt water)	Field	1	21
Sweet pepper	Powdery mildew	1.0 kg/ ha (100 gm/200 lt water)	Field	1-2	7
Tomatoes	Early blight	600 gm/ha (60 gm/200 lt water)	Field & green house	1-3	14
Soft wheat	Rust, powdery mildew, scald, scab.	500 gm/ha (50 gm/200 lt water)	Field	1-2	60
Wheat	Septoria spot.	1.0 kg/ ha (100 gm/200 lt water)	Field	1-2	60
Onion	Purple blotch	752 gm/ha (75 gm/200 lt water)	Field	1	60

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

• For more details about first aid & precautions please refer to first aid & precautions index.