

# TOVAP-30% EC

HERBICIDE  
EMULSIFIABLE CONCENTRATE

VAPCO

Class:  
Aryloxyphenoxypropionate<sup>(1)</sup>  
N.A.<sup>(2)</sup>

A selective systemic herbicide used for annual and narrow weeds such as: wild oats, canary grass and ryegrass in wheat fields.

**COMPOSITION:** Each Liter contains:

CLODINAFOP-PROPARGYL <sup>(1)</sup>	24% W/V (A.I)
CLOQUINTOCET-MEXYL <sup>(2)</sup>	6% W/V (A.I)

**PROPERTIES:**

**TOVAP-30% EC:** Selective systemic herbicide should be applied when weeds are actively growing.

**TOVAP-30% EC:** A herbicide that controls wild oats, canary grass and ryegrass in wheat fields.

**USES AND APPLICATION RATE:**

CROP	WEEDS	APPLICATION RATE / HECTARE	TIME OF APPLICATION
Wheat from 2 leaf to late jointing.	Wild oats	65 ml + 1 lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 5 leaf stage
		85 ml + 1 lt. of sticking materials / 100 lt. Water.	When weeds at 2 leaf to mid tillering stage
	Canary grass	85 ml + 1 lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 3 leaf stage
		125 ml + 1 lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 5 leaf stage
		160 ml + 1 lt. of sticking materials / 100 lt. Water.	When weeds at 2 leaf to early tillering stage
	Annual ryegrass	160 ml + 1 lt. of sticking materials / 100 lt. Water.	When weeds at 2 to 4 leaf stage
210 ml + 1 lt. of sticking materials / 100 lt. Water.		When weeds at 2 leaf to early tillering stage	

Donum = 1000 m<sup>2</sup>    Hectare = 10000 m<sup>2</sup>

**CAUTION:** Used only on the above mentioned crop.

• For more details about first aid & precautions please refer to first aid & precautions index.

**REMARKS:**

- Calibration of sprayer is important before application.
- Complete and homogeneous coverage while spraying is recommended.
- Do not apply if rainfall is expected within 2 hours.
- Do not apply to crops which are under stress. For example, due to very dry, very wet, frosty or diseased conditions, nutrient deficiency or high insect pressure.

**RE-ENTRY PERIOD:**

Do not enter the treated area without protective clothing until the drying of the spray solution

**GRAZING INSTRUCTIONS:**

Animals should be prevented from grazing in newly treated areas

**COMPATIBILITY:**

- It is preferable to make initial compatibility test before mixing with other herbicides before spraying.
- When **TOVAP-30% EC** is applied with compatible herbicides always use a recommended sticking product.
- The following herbicides are known to be incompatible with **TOVAP-30% EC**: Diuron, Triasulfuron, (Bromoxynil + Dicamba + MCPA) and all 2,4-D formulations. Apply these herbicides after 10 days at least from last treatment of **TOVAP-30% EC**, these compounds are sprayed separately.
- Can be compatible with insecticide like **ULTRACIDIN-40% EC**.

• For more details about first aid & precautions please refer to first aid & precautions index.