

VACOMIL-35% WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Acylalanine

A systemic fungicide effective in controlling fungal diseases such as: Blights, Downy mildew .

COMPOSITION: Each kg contains:

METALAXYL

35% W/W (A.I)

PROPERTIES:

VACOMIL-35% WP: A systemic fungicide effective in controlling fungal diseases on a wide range of crops.

VACOMIL-35% WP: Used as seed treatment to protect from soil diseases.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE / 20 LT. WATER	SAFETY PERIOD (DAYS)
Potato, Cucurbits	Downy Mildew, Late Blight	15 gm	7
Onion, Garlic	Downy Mildew, Blight, Mold	6 - 12 gm	7
Grapes	Downy Mildew	15 - 20 gm	7
Ornamentals	Downy Mildew	15 gm	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

NOTICE:

Used as seed dressing for the control of emergence diseases used at rate 200 gm product / 100 Kg seeds.

SLURRY TREATMENT:

Carefully mix the appropriate quantity of **VACOMIL-35% WP** for the corresponding quantity of seeds with sufficient water to obtain fluid mixture, for 3-5 minutes.

DRY TREATMENT:

Used a mechanical mixer, place the seed together with 0.5 – 1 lt. water/100 Kg seeds, shake for 1-2 minutes, then add the corresponding dose of **VACOMIL-35% WP**, and re-shake for about 3 minutes.

RE-ENTRY PERIOD:

48 hours from last application.

COMPATIBILITY:

Compatible with most other pesticide except alkaline compounds and compatibility test is recommended before application.

• For more details about first aid & precautions please refer to first aid & precautions index.