

VACOMIL-GOLD 68% WP

FUNGICIDE
WETTABLE POWDER

VAPCO

Class:
Acylalanine⁽¹⁾
Dithiocarbamate⁽²⁾

A special formula with a broad spectrum efficiency with systemic and contact action for the control of Downy Mildew, Blights and other diseases to help farmers to maximize their income.

COMPOSITION: Each kg contains:

METALAXYL-M (MEFENOXAM) ⁽¹⁾	4% W/W (A.I)
MANCOZEB ⁽²⁾	64 % W/W (A.I)

PROPERTIES:

VACOMIL-GOLD 68% WP: A Systemic fungicide which protects the treated plants efficiently.

VACOMIL-GOLD 68% WP: Has immediate effects against foliage and soil-borne diseases such as rots, blights, downy mildew diseases on vine, vegetables, sugar beet, barriers and pineapple.

VACOMIL-GOLD 68% WP: Not phytotoxic if used as recommended.

USES AND APPLICATION RATE:

CROP	DISEASE	APPLICATION RATE	NOTE	SAFETY PERIOD (DAYS)
Potatoes	Late and Early blight, Storage rots, Pink Rot	113 gm/donum in sufficient water to ensure thorough coverage of foliage (for aerial application), apply 113 gm in a minimum 50 Lt. water/donum.	Apply up to 3 applications/season, start application early-the first application should be applied before the leaves of the plants touch within a particular potato row. Second and third application is done 10-14 days.	14
Cucumber, Watermelon, Muskmelon, Squash	Downy mildew	127 gm/donum in sufficient water to obtain throughout coverage.	Begin applications when conditions are favorable for disease, but before infection, and continue at 14 days intervals until the threat of disease is over. The full rate of a protectant fungicide should be applied between VACOMIL-GOLD 68% WP spraying. avoid late season applications when plants reach full maturity or begin senescence.	5

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	DISEASE	APPLICATION RATE	NOTE	SAFETY PERIOD (DAYS)
Onion, Garlic (dry bulbs)	Downy Mildew	113 gm/donum in sufficient water to obtain thorough coverage	Apply up to 3 applications/season. Apply every 10-14 days.	7
Sugar beets	Downy Mildew	127 gm/donum	Begin applications when conditions are favorable for disease, but before infection, and continuo at 14 days intervals until the threat of disease is over.	7
Tomatoes	Late blight	127 gm/donum in sufficient water to obtain thorough coverage of ground application and in a minimum of 5.6 L of water/donum for aerial applications.	Begin applications when conditions are favorable for disease (before infection). Make up to 3 applications of VACOMIL-GOLD 68% WP at 14 days intervals. the full rate of a protectant fungicide should be applied between VACOMIL-GOLD 68% WP applications, apply the full rate of a protectant fungicide on a weekly schedule through reminder of the season.	5

Donum = 1000 m²

CAUTION:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hour from last application.

COMPATIBILITY:

Compatible with most pesticides, except alkaline materials, compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.