

VAPCOMIC-FLEX 18.5% SC

ACARICIDE/INSECTICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Avermectin⁽¹⁾
Nitroguanidine⁽²⁾

Acaricide and Insecticide contains two active ingredients (Thiamethoxam and Abamectin) with systemic, contact and ingestion action used to control piercing-sucking insects and mites that mentioned in the table of uses.

COMPOSITION: Each Liter contains:

ABAMECTIN⁽¹⁾
THIAMETHOXAM⁽²⁾

3.3% W/V (A.I)
15.2 % W/V (A.I)

PROPERTIES:

VAPCOMIC-FLEX 18.5% SC: Acaricide and Insecticide with systemic, contact and ingestion action.

VAPCOMIC-FLEX 18.5% SC: Contains Thiamethoxam and Abamectin used to control piercing-sucking insects and mites.

USES AND APPLICATION RATE:

CROPS	PESTS	APPLICATION RATE / 20 LT. OF WATER	APPLICATION TIME	SAFETY PERIOD (DAYS)
Citrus (lemon, orange, mandarina, pomelo, kumquat and grapefruit)	Aphids, Scales insects, Broad mite, Citrus black fly, Leafminer, Citrus rust mite, Thrips Fruit fly, Leafhopper, Mealybugs, Two spotted spider mite, Whiteflies	8-12 ml	- Do not use to nurseries. - Do not use during pre-bloom or bloom stage while bees are active. - Days interval between applications: *30 days at least for VAPCOMIC-FLEX 18.5% SC or any other product contains Abamectin. (applications must not exceed more 3 application per year), re-application can be done after *7 days at least for products containing Thiamethoxam.	7
Potato, Sweet potato, Artichoke, Ginger	Flea beetle (adult stage), leafminers Leafhopper, mites	7 ml	- Make the first application when pest appears. - Days interval between applications: 7 days at least. - Do not apply more than 2 applications/year.	14

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

- Used only on the above mentioned crops, and follow the recommended safety period before harvest.
- Do not use during pre-bloom or bloom stage in citrus while bees are active.
- Wait at least 5 days before placing beehives in the treated area.
- Spray **VAPCOMIC-FLEX 18.5% SC** after sunset and when temperature is below 12 °C.
- In potato, sweet potato, artichoke and ginger spray after 7 days at least (not more than 2 applications/year).
- In Citrus, spray **VAPCOMIC-FLEX 18.5% SC** or any other product contains Abamectin after 30 days at least. (Not more 3 application per year), for products containing Thiamethoxam after 7 days at least.
- Do not spray **VAPCOMIC-FLEX 18.5% SC** through any type of irrigation system.

RE-ENTRY PERIOD:

4 days from last application.

GRAZING:

Do not allow livestock to graze in treated area.

COMPATIBILITY:

Compatible with most pesticides except captan, compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.