

VAPCOMIC-PRO 50.4% SC

ACARICIDE/INSECTICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Avermectin⁽¹⁾
Carbazate⁽²⁾

Non Systemic Acaricide and Insecticide with contact action and long residual activity.

COMPOSITION: Each Liter contains:

ABAMECTIN⁽¹⁾
BIFENAZATE⁽²⁾

2.4% W/V (A.I)
48.0 % W/V (A.I)

PROPERTIES:

VAPCOMIC-PRO 50.4% SC: A unique combination of two active compounds (Bifenzate and Abamectin) that has an effect through contact action.

VAPCOMIC-PRO 50.4% SC: Combines of two active compounds with different mood of action to control all stages of mite and insect.

VAPCOMIC-PRO 50.4% SC: Used to control two-spotted spider mite, bud mite, European red mite, rust mite and leafminer.

USES AND APPLICATION RATE:

CROPS	PESTS	APPLICATION RATE / 20 LT. OF WATER	APPLICATION TIME	SAFETY PERIOD (DAYS)
Ornamentals	Two-spotted Spider mite, Bud mite, European Red Mite, Rust mite, Leafminers.	5 – 6 ml	- Apply when mites appear. - When sever infection, second application can be made after 7 days of first application. - Do not make more than two applications per season.	---
		9 ml		
Non-bearing fruit trees	Two-spotted Spider mite, Bud mite, European Red Mite, Rust Mite. Leafminers.	5 – 6 ml		---
		9 ml		
Nurseries, Forest trees, Conifer trees, Landscape trees	Two-spotted Spider mite, Bud mite, European Red mite, Rust mite. Leafminers.	5 – 6 ml	---	
		9 ml		

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

- Used only on the above mentioned crops, and follow the recommended safety period before harvest.
- Do not apply aerially, Do not apply through any type of irrigation system.
- Do not use in citrus nurseries.
- May be phytotoxic to apple fruit when mixed with captan.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Compatible with most pesticides except captan materials, compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.