

VAPCOMIC-1.8% EC

INSECTICIDE / ACARICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Avermectin

Insecticides and Acaricide with contact and stomach action effective to control insects in adult stage mentioned in table of use.

COMPOSITION: Each liter contains:

ABAMECTIN

1.8% W/V (A.I)

PROPERTIES:

VAPCOMIC-1.8% EC: Insecticide and Acaricide effective to control insects in Adult stage that are difficult to control like: leafminer, psylla and Red Spider Mite.

VAPCOMIC-1.8% EC: Highly active for a long period due to its content of a high percent of the active isomer Avermectin B1a.

VAPCOMIC-1.8% EC: Used on economic crops like: Tomato, Cucumber, Strawberry, Citrus and Ornamentals.

USES AND APPLICATION RATE:

CROP	PEST	APPLICATION RATE / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomatoes	Leafminers, Red Spider mites.	10 - 20 ml	7
Cucumber	Leafminers, Red Spider mites.	10 - 20 ml	3
Strawberries	Leafminers, Red Spider mites.	10 - 20 ml	7
Orange, Pomelo, Lemon, Grape-fruit, Mandarin, Kumquat	psylla, rust mite.	10 ml	10
Ornamentals	Leafminers, Red Spider mites.	10 - 20 ml	---

CAUTION:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

REMARKS:

- * Do not spray the crop more than 5 times during one season.
- * Results of control appears after 3-5 days of spraying, however effect of the product appears immediately after spraying.
- * Quantity of the solution must not exceed 110 ml of **VAPCOMIC 1.8% EC** /1000 m² for vegetables and ornamentals.

• For more details about first aid & precautions please refer to first aid & precautions index.

RE-ENTRY PERIOD:

24 hours from last application, the spray should be allowed to dry on the foliage before re-entry into the treated areas.

PHYTOTOXICITY:

Non-phytotoxic, but do not spray it on carnations and sedges, toxic to Pome fruits when mixed with Captan (fungicide) containing compounds.

COMPATIBILITY:

Compatible with most pesticides, compatibility test is recommended before spraying.

*** OTHER TRADE NAMES: ABAMEX-1.8% EC, BAMECTIN-1.8% EC, MAYSAMECTIN-1.8% EC, VAPCOMIC-2% EC**

• For more details about first aid & precautions please refer to first aid & precautions index.