

# B-BROVAP 10%

Fertilizer  
Soluble Liquid

VAPCO

Class:  
Trace elements

B-BROVAP 10% Fertilizer contain boron which used to control boron deficiency in the crops in the table of use

COMPOSITION: Each liter contains:

Boron 10% W/W Equivalent to 13.5% W/V (A.I)

## PROPERTIES:

**B-BROVAP 10%:** Is an effective readily available source of boron for soil and foliar applications to plants

**Foliar Applications:** Boron is not easily translocated within the plant, therefore multiple foliar applications at low rates during the growing season are most effective.

**Soil Applications :** Soil applied boron must be moved into the root zone to be absorbed by the plant. This can be accomplished by irrigation, rainfall, or tillage.

## USES AND APPLICATION RATE :

CROP	FOLIAR/ 20 L WATER	SOIL/DONUM
	APPLY IN SUFFICIENT WATER FOR THOROUGH COVERAGE	APPLY IN FLUID COVERAGE FERTILIZER OR WATER
Apples, Apricots, Avocados, Olives, Pistachios, Strawberries	48-95 ml	1-2.85 L
Almonds, Filberts	24-71 ml	1-2 L
Caneberries, Grapes, Melons, Walnuts	95-237 ml	1-2 L
Citrus Fruits, Cherries, Pears, Peaches, Plums	95-189 ml	1-2 L
Alfalfa	95-189 ml	1-2.85 L
Beets (Sugar)	95-189 ml	1-2.85 L
Cotton, Flax, Mint	48-95 ml	0.5-1 L
Corn	48-95 ml	1-2 L
Peanuts	48-95 ml	0.5-1 L
Soybeans, Sugarcane	48-95 ml	1-2 L
Sunflowers, Wheat	48-95 ml	0.5-1 L
Beets, Carrots, Cucumbers, Lettuce, Onions, Radishes, Spinach, Turnips	48-95 ml	1-2 L
Broccoli, Cabbage, Cauliflower	95-189 ml	1-2.85 L

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	FOLIAR/ 20 L WATER	SOIL/DONUM
	APPLY IN SUFFICIENT WATER FOR THOROUGH COVERAGE	APPLY IN FLUID COVERAGE FERTILIZER OR WATER
Celery	95-189 ml	1-2.85 L
Peppers, Potatoes, Tomatoes	48-95 ml	1-2.85 L
Beans, Lentils, Peas	48-95 ml	1-2.85 L
Turf and Ornamental	47-142 ml	----
Other Crops	Apply as early in the growing season as possible, providing there is adequate leaf coverage	

**CAUTION:**

- Over application can cause foliar damage or serious crop injury.
- Drip Irrigation: The required amount of this product should be applied over the growing season with multiple applications 1 to 2 weeks apart.
- Do not use in soil applications where it will directly contact the seed.

**COMPATIBILITY:**

Compatible with many liquid fertilizers, micronutrients, and pesticides, compatibility test is recommended before spraying.