

FENATODE-10% GR

NEMATICIDE
GRANULES

VAPCO

Class:
Phosphoramidate

Systemic nematicide with contact action used to control nematodes in tomato, citrus, banana, and potato.

COMPOSITION: Each kg contains:

FENAMIPHOS

10% W/W (A.I)

PROPERTIES:

FENATODE-10% GR: Highly effective Organophosphorus nematicide with cholinesterase inhibitor activity to control nematodes.

FENATODE-10% GR: Systemic nematicide with contact action, absorbed by the roots, and translocated to the leaves protect the plants from new injuries.

FENATODE-10% GR: Readily distributed in the soil so that it controls Nematodes present in deep soil layers. Also It has a residual activity for several months.

FENATODE-10% GR: Stimulates growth, increase yield and improve its quality.

FENATODE-10% GR: Effective against soil pests like populations of ectoparasitic, endoparasitic, free-living, cyst-forming and root-knot nematodes.

USES AND APPLICATION RATE:

Use **FENATODE-10% GR** only in wet soil either during winter season or after irrigation. Mix the granules mildly with soil immediately after application.

CROP	PEST	DOSAGE	SAFETY PERIOD (DAYS)
Citrus * Small trees * Old trees	Slow Decline Nematode	Used once in spring in rate 40 gm / m ² . Add 20 - 40 gm / m ² after harvest immediately, in case of sever infestation apply again with the same rate in autumn by applying under trees and mix to depth 5 -10 cm and irrigate in rate 25 lt. / m ² .	90
Tomatoes	Nematode	In open filed apply on rows before 14 days before seedling plant in the soil, soil raking to depth 5 - 10 cm then irrigate heavily 25 lt. water/m ² in rate of 1.5 - 3 kg./donum. In green house apply 14 days before planting in the rate of 4.5 kg/donum, soil raking to depth 5-10 cm and irrigate heavily.	90
Potatoes	Nematode	Apply in the row with tubers then cover the row and irrigate heavily in a rate of 1.5 kg/donum. To treat the whole field apply in rate 4 - 5 kg/donum two weeks before planting tubers.	60
Banana	Nematode	New shoots: cover shoots with mixture of clay and water in addition to FENATODE-10% GR in the rate of 8 kg/100 lt. clay, retreat after 4 months in case of sever infection by applying granules 25 - 30 cm away from the hole during boot stage then irrigate heavily.	60

Donum = 1000 m²

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on the above mentioned crops and follow the recommended safety period before harvest.

COMPATIBILITY:

Compatible with the same formulation pesticide, incompatible with alkaline materials, compatibility test is recommended before 24 hours from application

• For more details about first aid & precautions please refer to first aid & precautions index.