

AGRINATE-24% SL

INSECTICIDE
SOLUBLE CONCENTRATE

VAPCO

Class:
Oxime Carbamate

Systemic insecticide used to control a wide range of insects on agricultural crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

METHOMYL

24% W/V (A.I)

PROPERTIES:

AGRINATE-24% SL: Has a fast knockdown on a wide range of insects.

AGRINATE-24% SL: Has a broad spectrum activity against a wide range of insects at all developing stages (eggs, larvae).

AGRINATE-24% SL: Is extremely active against the most difficult insect pests such as borer, fruitworm and leafminer and against insects which have become resistant to Organophosphate and Pyrethroid compounds.

AGRINATE-24% SL: Quickly decomposes in nature, which decreases the safety period for vegetables and fruit trees.

AGRINATE-24% SL: Has no effect to the environment due to it's rapid decomposition.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomatoes	Fruitworm, Leafworm	11-15	7
Cabbage, Cauliflower	Green Worms, Leafworm, Aphids, Thrips.	11-15	7
Cucumber, Squash, Watermelon, Beans, Jew's mallow, Pepper, Eggplant, Potatoes.	Leafworm, Aphids, Thrips.	15	7
Apples, Pear, Peach, Apricot, Cherries, Almonds, Citrus	Green And Black Aphids, Leafminer, Stem Borer, Psylla, Leafroller, Thrips, Leafhopper.	10-15	7
Grapes	Grape Berry moth, Leafhopper.	11-15	7
Corn	Cutworm, Corn Borer, Aphids, Beetles.	15	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Ornamentals (Open Field, Green Houses)	Leafroller, Leafminers, Bugs, Leafworm, Leafhopper, Aphids, Thrips.	11-15	---

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

72 hours after the last application.

REMARKS:

- Use ground sprayers with low volume pressure, use for large farms and it is not recommended for use in gardens.
- Fill 1/4 or 1/2 of the spray tank with water, then add the recommended dose of **AGRINATE-24% SL** stir, fill the tank and start spraying.
- Use the recommended dose at the beginning of infestation, repeat spraying 5-7 days intervals if needed.
- In case there is more than one dose recommended, use the lower dosage for young crops or when the injury is slight.

COMPATIBILITY:

Compatible with most common insecticides and fungicides, but compatibility test is recommended before spraying.

*** OTHER FORMULATIONS: AGRINATE-25% (SP,WP), AGRINATE-90% SP**

• For more details about first aid & precautions please refer to first aid & precautions index.