

# AGRINATE-25% SP

INSECTICIDE  
SOLUBLE POWDER

VAPCO

Class:  
Oxime Carbamate

Systemic insecticide used to control a wide range of insect pests which attack agricultural crops.

**COMPOSITION:** Each kg contains:

**METHOMYL**

**25% W/W (A.I)**

## PROPERTIES:

**AGRINATE-25% SP:** Packed in water soluble sachets.

**AGRINATE-25% SP:** Has a fast knockdown on a wide range of insects.

**AGRINATE-25% SP:** Has a broad spectrum activity against a wide range of insects at all developing stages (eggs, larvae).

**AGRINATE-25% SP:** Quickly decomposes in nature, which decreases the safety period for Vegetables and Fruit trees.

**AGRINATE-25% SP:** Has no effect to the environment due to it's rapid decomposition.

**AGRINATE-24% SL:** Has an emetic material and it's unique from other products by its blue color.

## USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE	SAFETY PERIOD (DAYS)
Tomatoes	Fruitworm, Leafworm	2 - 4 bags/drum	7
Cabbage, Cauliflower	Green worms, Leafworm, Aphids, Thrips.	2 - 4 bags/drum	7
Cucumber, Squash, Watermelon, Beans, Jew's mallow, Pepper, Eggplant, Potatoes.	Leafworm, Aphids, Thrips.	4 bags/drum	7
Corn	Cutworm, Corn borer, Aphids, Beetles.	4 bags/drum	7
Apples, Pear, Peach, Apricot, Cherries, Almonds and Citrus	Green and black aphids, Leafminer, Stem borer, Psylla, Leaf roller, Thrips, Leafhopper.	4 bags/drum	14
Grapes	Grape Berry moth, Leafhopper.	4 bags/drum	7
Ornamentals (Open Field, Green Houses)	Cotton Leafworm, Cutworm, Bugs, Aphids, Thrips, Leafhopper.	4 bags/drum	--

Drum = 200 lt. of water.

Bag = 200 gm of **AGRINATE-25% SP**

• For more details about first aid & precautions please refer to first aid & precautions index.

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

72 hours from last application.

**REMARKS:**

- Use ground sprayers with low volume pressure for large farms and it is not recommended for use in gardens.
- Fill 1/4 or 1/2 of the spray tank with water, then add the recommended dose of **AGRINATE-25% SP** stir, fill the tank and start spraying.
- Use the recommended dose at the beginning of infestation, repeat spraying 5 - 7 days intervals.
- In case there is more than one dose recommended, use the lower dosage for young crops or when the injury is slight.

**COMPATIBILITY:**

Compatible with most common insecticides and fungicides, but compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.