

AZARIL-50% WP

PUBLIC HEALTH INSECTICIDE
WETTABLE POWDER

VAPCO

Class:
Organophosphate

An organophosphorus insecticide, with contact and stomach action effective to control public health insects mentioned in the table.

COMPOSITION: Each kg contains:

AZAMETHIPHOS

50% W/W (A.I)

PROPERTIES:

AZARIL-50% WP: An organophosphorus insecticide with contact and stomach action, to control ants, flies, cockroaches, spiders and fleas in household and other institutional where insects can be exist.

AZARIL-50% WP: Gives rapid knockdown, and has long residual activity.

AZARIL-50% WP: A formulation that is suitable for application by spraying or paint on walls.

USES AND APPLICATION RATE:

SITUATION	PESTS	APPLICATION RATE / 20 LT. WATER	NOTES
- Empty dairies, stables, animals hangars, empty abattoirs. - Kitchens, restaurants, canteens, empty hospitals. - Empty bakeries, warehouses and other commercial premises.	Cockroaches, Spiders, Flies, Ants and Fleas	100 – 200 gm	- Spray where insects enter or congregate. - Cockroaches hiding places such as under benches, in cupboards, behind sinks, stoves and refrigerators. - Spray near entrances with paying particular attention to cracks and crevices. - Protect food utensils from contamination.
- Poultry sheds and egg packing establishments	Flies	10 gm of AZARIL-50% WP + 40 gm sugar, added to 40 ml water to make up a paintable mixture	Paint strips of 5x30 cm where flies congregate, e.g. warm wall areas partitions, milk pipes, etc. 250 gm of AZARIL-50% WP / sugar mixture in 200 ml of water is sufficient to treat a building of approx. 100 m ² ground area and 200 m ² wall and ceiling surfaces.
Garbage disposal and refuse area	Flies	50 gm	For knockdown Fly control in garbage tips apply a minimum of 50 liters spray mixture / hectare. Spray or mist directly on to rubbish tips.

RE-ENTRY PERIOD: 24 hours from the last treatment.

Hectare = 10000m²

COMPATIBILITY:

Compatible with most pesticides except alkaline ones and its preferable to make compatibility test 24 hours before application.

RESIDUAL ACTIVITY: About 8 Weeks.

• For more details about first aid & precautions please refer to first aid & precautions index.