

CINTURA-48% SC

Insecticide
Suspension Concentrate

VAPCO

Class:
Diamide

Insecticide to control of various lepidopteran pests in crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

FLUBENDIAMIDE

48% W/V (A.I)

PROPERTIES:

CINTURA-48% SC: Systemic insecticide active primarily by ingestion.

CINTURA-48% SC: It works to activate of the ryanodine receptors in the calcium release channel, leading to contraction and paralysis of the insects.

USES AND APPLICATION RATE :

CROP	PEST	APPLICATION RATE	PHI
Broccoli, Cabbage, Cauliflower	Diamondback moth, cabbage white butterfly, cluster caterpillar, heliothis	75 -100 ml / ha	3 DAYS
Cucumbers, Melons, Pumpkin, Squash.	Heliothis,	100 ml / ha	3 DAYS
Eggplant, Peppers, Tomato	Heliothis, tomato leafminer	100 -150 ml / ha	3 DAYS
Lettuce, Chard, Rocket, Celery.	Heliothis	75 -100 ml / ha	3 DAYS
Potato, Sweet Potato, Beetroot, Carrot, Radish, Turnip, Dasheen (Taro).	Diamondback moth, cabbage white butterfly, cluster caterpillar, heliothis	75 -100 ml / ha	3 DAYS
	tomato leafminer	100 -150 ml / ha	
Strawberry	cluster caterpillar, heliothis	100 ml / ha	3 DAYS

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

- Used only on the above-mentioned crops and follow the recommended safety period before harvest.
- Application of this product to crops other than those specified on this label may cause symptoms of phytotoxicity.
- A maximum of three applications may be applied to one crop.
- spray intervals of 7 - 14 days. Use the shorter application interval (7 to 10 days) during periods of high pest pressure or rapid crop growth.
- Use the higher application rate during high pest pressure.
- START applications once local economic spray thresholds are reached. Apply at egg hatch in crops with nil thresholds for damage. Otherwise, apply at egg hatch or very soon after egg hatch to target young larvae.

COMPATIBILITY:

Compatible with most pesticides and its preferable to make compatibility test before spraying.

RE ENTRY PERIOD:

Do not enter into treated areas until the spray has dried