CYPERMETHRIN-5% EC

INSECTICIDE EMULSIFIABLE CONCENTRATE



Non-systemic insecticide with contact and stomach action, used to control many of insect pests which attack agricultural crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

CYPERMETHRIN

5% W/V (A.I)

PROPERTIES:

 $\textbf{CYPERMETHRIN-5} \% \ \textbf{EC:} \ \textbf{Non-systemic pyrethroidal insecticide with a contact and stomach}$

action for the control of insects mentioned in the table of uses. **CYPERMETHRIN-5% EC:** Very effective insecticide with no residue in the environment.

CYPERMETHRIN-5% EC: Has a residual effect on the treated crops.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	PERIOD
Tomato, Pepper, Eggplant, Okra, Potato	Cutworm, Whitefly Cotton Leafworm.	10-20 10-12	7 7
Apple, Pear	Caterpillars, Borers	10-20	7
Cotton	Cutworm, Cotton Leafworm. Aphids, Leaf Rollers.	20-40 10-20	7 7
Grapes	Grapes berry Moth	10-20	7

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours from last application.

COMPATIBILITY:

Compatible with most insecticides and fungicides except alkaline compounds, but compatibility test is recommended before spraying.

[•] For more details about first aid & precautions please refer to first aid & precautions index.