

# CYPOTHRIN-25%EC

INSECTICIDE  
EMULSIFIABLE CONCENTRATE

VAPCO

Class:  
Pyrethroid

A non-systemic pyrethroidal insecticide with a contact and stomach action, effective against wide range of insect pests attacking agricultural crops that are mentioned in the table of uses

**COMPOSITION:** Each litre contains:

**CYPERMETHRIN**

**25% W/V (A.I)**

**PROPERTIES:**

**CYPOTHRIN-25% EC:** A pyrethroidal insecticide, moderately toxic to human and animals, in conditions of use.

**CYPOTHRIN-25% EC:** Extremely active against cutworms, leafhoppers, flea beetles, cucumber beetles, aphid, whiteflies, caterpillar, thrips and other pests on broccoli, cauliflower, cabbage, lettuce, garlic, leeks, and onion.

**CYPOTHRIN-25% EC:** Non phytotoxic when used as recommended.

**USES AND APPLICATION RATE:**

CROP	PESTS	APPLICATION RATE ML / 20L WATER	SAFETY PERIOD (DAYS)
Broccoli, Cabbage, Cauliflowe, Lettuce.	Cutworms, Caterpillar, Leafhoppers, Flea beetles, Cucumber beetles, Aphid, Whiteflies, Diamond back moth, Budworms	5 - 10	7
	Worms, Thrips	7 - 10	
Garlic, Leeks, Onion	Thrips, Aphid, Worms, Bugs	7 - 10	7

**CAUTION:**

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Lower rate of **CYPOTHRIN-25% EC** should be used under light and moderate insect population
- Higher rates of **CYPOTHRIN-25% EC** should be used to control heavy insect populations.
- The interval days between one spray and the next should be not less than 7 day.

**RE-ENTRY PERIOD:**

48 hours from last application.

**COMPATIBILITY:**

Compatible with most pesticides, but compatibility test is recommended before spraying.

- For more details about first aid & precautions please refer to first aid & precautions index.