

# DESTROYER-48% SC

INSECTICIDE  
SUSPENSION CONCENTRATE

VAPCO

Class:  
Cyanoimidamide

Systemic insecticide to control sucking and some chewing insects on a wide range of agricultural crops.

**COMPOSITION:** Each liter contains:

**THIACLOPRID**

**48% W/V (A.I)**

**PROPERTIES:**

**DESTROYER-48% SC:** A new generation insecticide belongs to Neonicotinoid family.

**DESTROYER-48% SC:** Acute contact and stomach poison, with systemic properties.

**DESTROYER-48% SC:** Stable to different environment conditions such as rain and sunlight which enable plants to absorb the pesticide in effective way.

**DESTROYER-48% SC:** For use by foliar application against sucking and chewing insects on a wide range of agricultural crops.

**DESTROYER-48% SC:** Controls insects which are resistant to other insecticides.

**DESTROYER-48% SC:** Has no effect on bees and natural enemies making it a suitable choice to be used in IPM programs.

**USES AND APPLICATION RATE:**

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Tomatoes, Cucumber, Pepper, Watermelon, Muskmelon.	Aphids, Whitefly	4-6	3
Apple, Pear,	Aphids, Furry rose beetle Leafhopper Leaf Borers, Codling Moth Green Apple Bug Mealybug Pear Psylla	4-5 6 4-5 4-5 1.25-2 2-4	14
Peach, Apricot, Nectarine, Almond	Aphids	2-4	14
Grapes	Leafhopper	3	14

• For more details about first aid & precautions please refer to first aid & precautions index.

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

48 hours from last application.

**COMPATIBILITY:**

Compatible with all pesticides except alkaline and acidic compounds, but compatibility test is recommended before spraying.

**\* OTHER TRADE NAME: DECLOPRID-48% SC**

• For more details about first aid & precautions please refer to first aid & precautions index.