DOUBLE ACT

PUBLIC HEALTH INSECTICIDE GEL BAIT



Cockroach gel bait to control cockroaches in the situations specified in the table of uses.

COMPOSITION: Each KG Contains:

INDOXACARB 0.06% W/W (A.I)

PROPERTIES:

DOUBLE ACT: Control pest species of cockroaches including American and German.

DOUBLE ACT: Cockroach gel can be used indoors or outdoors.

DOUBLE ACT: The mode of action of the active ingredient indoxacarb allows cockroaches to

consume the bait and return to the harbourage site and to contaminate other

cockroaches which lead to reduction in the population of cockroach

USES AND APPLICATION RATE:

SITUATION	PESTS	APPLICATION RATE	REMARKS
Domestic, commercial and public service buildings including factories, farm buildings, food processing establishments, homes, hospitals, kitchens, offices, storerooms, restaurants, retail outlets Vehicles, Other situations.	American Cockroach German Cockroach other common cockroaches	Apply at least 2 spots/m2. For heavy infestations increasing the number of spots (up to 5/m2) maybe required or if American Cockroaches are the main pest. Recommended spot size is approximately 5 mm diameter (0.1 g).	Can be used indoor and outdoor. Indoor applications can include cracks and crevices along walls or floors, behind or under equipment, under tables, within framing of tables or furniture, around sinks, within or near garbage collection areas, attics, crawl space and cracks and crevices in cabinets. Apply where Cockroach are existed in the hidden areas, between construction elements or other cracks and crevices. DOUBLE ACT should be applied as a small spot to areas not able to be contacted by children or pets. Several small applications distributed within an area are more effective than 1 or 2 applications of large quantities. In some cases a small thin bead (less than 2.5 mm wide and 5 cm long) may be appropriate, Inspect Bait placements periodically.

[·] For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

- Use the product only as the above-mentioned recommendation and doses.
- Avoid applying to surfaces recently treated with residual sprays and surfaces which treated with double act should not be treated with residual sprays.