

# EL-TORO 25% SC

INSECTICIDE  
SUSPENSION CONCENTRATE

VAPCO

Class:  
Nitroguanidine

Highly effective Systemic Insecticide with contact and stomach action, used to control many of sucking and chewing insects on a wide range of crops listed in the table of uses.

**COMPOSITION:** Each liter contains:

**THIAMETHOXAM**

**25% W/W (A.I)**

**PROPERTIES:**

**EL- TORO 25% SC** : Highly effective systemic insecticide with contact and stomach action.

**EL- TORO 25% SC** : Highly effective with low dosage and long lasting effect.

**EL- TORO 25% SC** : Has a special formula that interferes with information transfer between nerve cells making the insects become paralyzed.

**USES AND APPLICATION RATE:**

| CROP   | PESTS                           | APPLICATION RATE ML / 20 LT. WATER | SAFETY PERIOD (DAYS) |
|--|---------------------------------|------------------------------------|----------------------|
| Cucumber, Muskmelon, Pumpkin, Squash, Watermelon | Aphids                          | 2 – 4                              | 7                    |
|  | Beetles, Leafminer, Whiteflies. | 4 – 8                              |                      |
| Eggplant, Pepper, Tomato                         | Aphids, Leafhoppers.            | 3 – 4                              | 7                    |
|  | Whiteflies, Weevil, Bugs.       | 4 – 8                              |                      |
| Lettuce, Celery, Parsley, Spinach                | Aphids, Beetles, Leafhoppers    | 2 – 4                              | 7                    |
|  | Whiteflies                      | 4 – 8                              |                      |
| Cabbage, Cauliflower                             | Aphids, Beetles                 | 2 – 4                              | 7                    |
|  | Thrips, Whiteflies              | 4 – 8                              |                      |
| Mint   | Aphids, Beetles, Leafhoppers    | 2 – 4                              | 7                    |
| Carrot, Beet, Turnip                             | Aphids, Beetles, Leafhoppers    | 2 – 4                              | 7                    |
|  | Whiteflies                      | 4 – 6                              |                      |
| Radish   | Aphids, Beetles, Leafhoppers    | 2 – 4                              | 7                    |
|  | Whiteflies                      | 4 – 6                              |                      |
| Potato, Sweet Potato                             | Aphids                          | 4                                  | 14                   |
| Barley   | Aphids                          | 6                                  | 21                   |

• For more details about first aid & precautions please refer to first aid & precautions index.

| <b>CROP</b>   | <b>PESTS</b>  | <b>APPLICATION RATE ML / 20 LT. WATER</b> | <b>SAFETY PERIOD (DAYS)</b> |
|---|---|---|-----------------------------|
| <b>Lemon, Orange, Pummelo, Mandarin, Grapefruit</b> | Aphids, Leafhoppers   | 4 – 6                                     | 14                          |
|   | Psyllid, Citrus Fly, Leafminer, Mealybugs, Scales insects, Whiteflies | 6 – 8                                     |                             |
|   | Fruit Fly, Weevils, Plant Bugs, Borers, Thrips.                       | 7 – 8                                     |                             |
| <b>Apples, Loquat, Quince. Pear.</b>                | (post-bloom): Leafhoppers   | 3 – 4                                     | 14                          |
| <b>Apricot, Peach, Nectarine, Plum</b>              | Leafhoppers   | 3 – 4                                     | 14                          |
|   | Aphids  | 4 – 6                                     |                             |
|   | Fruit Fly, Bugs, Thrips   | 7 – 8                                     |                             |
| <b>Grape</b>  | Beetle, Leafhoppers, Mealybugs  | 2 – 5                                     | 5                           |
| <b>Strawberry</b>                                   | Aphids, Leafhoppers   | 2 – 4                                     | 3                           |
|   | Bugs, Fruit weevil  | 6   |                             |

**CAUTION:**

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- Do not spray during flowering stage.

**RE-ENTRY PERIOD:**

24 hours from last application.

**COMPATIBILITY:**

Compatible with most pesticide except highly alkaline and oxidative materials, but compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.