

EMAVAP-2% EC

INSECTICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Avermectin

Non-systemic insecticide which penetrates leaf tissues by translaminar movement, stop feeding and subsequently die after 2-4 days of lepidopteran insects

COMPOSITION: Each liter contains:

EMAMECTIN BENZOATE

2% W/V (A.I)

PROPERTIES:

EMAVAP-2% EC: Non-systemic insecticide for the control of worms and moths attacking a wide spectrum crops, it is active against insects which have become resistant to other traditional insecticides.

EMAVAP-2% EC: Penetrates leaf tissues, which mean a long lasting residual and killing effect.

EMAVAP-2% EC: Non-phytotoxic to plants.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	COMMENTS	SAFETY PERIOD (DAYS)
Cabbage, Broccoli, Cauliflower	Beet armyworm, Diamond back moth, Cabbage moth. Cabbage leaf roller, Leafminer.	4.25 - 8.5	Apply when larvae are first observed. Additional applications may be repeated to maintain control.	14
		5.6 - 8.5		
Lettuce, Celery, Spinach	Beet armyworm, Tobacco budworm. Cabbage leaf roller, Leafminer.	4.25 - 8.5		7
		5.6 - 8.5		
Eggplant, Pepper, Tomato	Beet armyworm, Cabbage leaf roller, Tobacco budworm, Tomato Fruit worm. Alfalfa looper, Leafminer	4.25 - 8.5	7	
		5.6 - 8.5		
Apple, Pear, Quince	Bud worms, Leafminers, Leaf rollers, mites	4.25 - 8.5	14	

• For more details about first aid & precautions please refer to first aid & precautions index.

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

48 hours from last application.

COMPATIBILITY:

Is compatible with most insecticides, fungicides, and foliar fertilizers, but compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.