

# EMAVAP-5% SG

**INSECTICIDE  
SOLUBLE GRANULES**

**VAPCO**

Class:  
**Avermectin**

**Non-systemic insecticide which penetrates leaf tissues by translaminar movement, stop feeding and subsequently die after 2-4 days of lepidopteran insects**

**COMPOSITION:** Each kg contains:

**EMAMECTIN BENZOATE**

**5% W/W (A.I)**

**PROPERTIES:**

**EMAVAP-5% SG:** Non-systemic insecticide penetrates leaf tissue, acts by contact action for the control of moths and worms on crops mentioned in the table of uses.

**USES AND APPLICATION RATE:**

<b>CROP</b>	<b>PESTS</b>	<b>APPLICATION RATE ML / 20L WATER</b>	<b>COMMENTS</b>	<b>SAFETY PERIOD (DAYS)</b>
<b>Cabbage, Broccoli, Cauliflower.</b>	Beet armyworm, Diamond back moth, Cabbage white butterfly, Leafminers.	3.5 – 7.0	Apply when larvae are first observed. Additional Applications may be repeated to maintain control.	14
				7
<b>Lettuce, Celery, spinach.</b>	Cutworms	3.5 – 7.0		7
<b>Eggplant, pepper, Tomato.</b>	Cutworms, Cabbage looper, Tobacco budworm, Tomato Fruit worm, Leafminers.	3.5 – 7.0		7
<b>Apple, pear, quince.</b>	Bud worms, Leafminers, Cutworms, Mites.	3.5 – 7.0		14

**CAUTION:**

Used only on above mentioned crops and follow the recommended safety period before harvest.

**RE-ENTRY PERIOD:**

24 hours from last application.

**COMPATIBILITY:**

Is compatible with most insecticides, fungicides, and foliar fertilizers, but compatibility test is recommended before spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.