

ENYO-10% SC

INSECTICIDE
SUSPENSION CONCENTRATE

VAPCO

Class:
Tetramic Acid

A systemic insecticide acting mainly by ingestion and also by contact, affecting eggs and larvae, with additional effects on adult fecundity, used to control various insect pests on vegetable, fruit tree and other crops mentioned in the table of uses.

COMPOSITION: Each liter contains:

SPIROTETRAMAT

10% W/V (A.I)

PROPERTIES:

ENYO-10% SC: Is fully systemic, moving through phloem and xylem to all plant tissues including new shoot, leaf and root growth.

ENYO-10% SC: Is active primarily by ingestion against immature target pest life stages, with additional effect on female adult fecundity used to control various insect pest in vegetables mentioned in the table of uses.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20L WATER	DAYS INTERVAL BETWEEN APPLICATIONS	RECOMMENDATIONS	SAFETY PERIOD (DAYS)
Grapefruit, Lemon, Mandarin, Orange, Pummelo.	Aphids, Psyllid, Scale insect, Thrips, Whiteflies, Bugs.	12-15	21	- Do not apply this product within 10 days prior to bloom, during bloom, or until petal fall is complete. - Apply with spreading agent in tank mix except for grapefruit. -Maximum ENYO-10% SC allowable per crop season 71 ml.	3
Grape, Kiwifruit	Mealybugs, Grape, phylloxera, whiteflies.	7-12	30	- When mixing with adjuvant, it is recommended that a high quality spreading/penetrating adjuvant be used and make a small scale trails. -Maximum ENYO-10% SC allowable per crop season 44.4 ml.	7
Apple, Loquat, Pear, Quince.	Aphids, Pear psylla, Mealybug, Whiteflies.	9-13	14	-Do not apply until petal fall. -Maximum ENYO-10% SC allowable per crop season 88.7 ml.	7
Apricot, Cherry, Plum, Prune.	Aphids, Mealybug, Whiteflies, Pear Psylla.	9-13	14	-Do not apply until petal fall. -Maximum ENYO-10% SC allowable per crop season 88.7 ml.	7

• For more details about first aid & precautions please refer to first aid & precautions index.

CROP	PESTS	APPLICATION RATE ML / 20L WATER	DAYS INTERVAL BETWEEN APPLICATIONS	RECOMMENDATIONS	SAFETY PERIOD (DAYS)
Almond, Cashew, Chestnut, Hazelnut, Pistachio, Walnut.	Aphids, Mealybug, Phylloxera, Whiteflies	9-13	14	-Do not apply until petal fall. -Maximum ENYO-10% SC allowable per crop season 76.3 ml.	7
Avocado, Guava, Mango, Papaya.	Aphids, Scale insects, Mealybug, Whiteflies.	12-15	14	-Maximum ENYO-10% SC allowable per crop season 88.7 ml.	3
Okra, Eggplant, Pepper, Tomato.	Aphids, Psyllids, Whiteflies.	4-7	7	-Maximum ENYO-10% SC allowable per crop season 35.5 ml.	3
Celery, Lettuce, Parsley, Spinach.	Cutworms.	4-7	7	-Maximum ENYO-10% SC allowable per crop season 35.5 ml.	3
Bean, Pea, Lupin, , Cowpea, Chickpea, Lentil.	Aphids, Whiteflies.	4-7	7	-Maximum ENYO-10% SC allowable per crop season 35.5 ml.	3 for fresh and edible legumes 7 for dry legumes
Broccoli, Cabbage, Cauliflower.	Aphids, Whiteflies.	4-7	7	-When mixing with adjuvant, it is recommended that a high quality spreading/penetrating adjuvant be used and make a small scale trails.. -Maximum ENYO-10% SC allowable per crop season 35.5 ml.	3
Artichoke, Ginger, Potato, Sweet Potato, Turmeric.	Aphids, Psyllids, Whiteflies	6-7	7	-Maximum ENYO-10% SC allowable per crop season 35.5 ml.	7

CAUTION:

- Used only on above mentioned crops and follow the recommended safety period before harvest.
- It is not recommended to add **ENYO-10% SC** to the tank mix with spray oil with captan and/or sulfur.
- Do not use the pesticide during the period of petals falling on fruit trees.
- Do not apply in plastic or glass houses.
- Apply when pest first appear and before pest population increase.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Compatible with most pesticides, but compatibility test is recommended before spraying.

- For more details about first aid & precautions please refer to first aid & precautions index.