

FENIKILL-30% EC

INSECTICIDE
EMULSIFIABLE CONCENTRATE

VAPCO

Class:
Organophosphate⁽¹⁾
Pyrethroid⁽²⁾

A special combination used to control various insect pests.

COMPOSITION: Each liter contains:

FENITROTHION⁽¹⁾
FENVALERATE⁽²⁾

25% W/V (A.I)
5% W/V (A.I)

PROPERTIES:

FENIKILL - 30% EC: Non-systemic insecticide with a contact and stomach action for the control of insects mentioned in the table of uses.

FENIKILL - 30% EC: Not phytotoxic if used as recommended but it may cause injuries for Cruciferous and some fruit trees if the dosage was high.

USES AND APPLICATION RATE:

CROP	PESTS	APPLICATION RATE ML / 20 LT. WATER	SAFETY PERIOD (DAYS)
Cabbage, Cauliflower, Potato, Tomato, Onion, Garlic, Squash, Beans, Muskmelon, Watermelon.	Caterpillars, Beetles, Thrips, Aphids	30 - 45	15
Grape, Citrus, Apple, Pear, Peach, Plum, Apricot, Cherries, Nectarine and Almond.	Codling moth, Fruit Moth, Fruit flies, Caterpillars, Beetles, Grape berry moth, Leafminers, Mealy bugs.	40 - 50	15
Wheat	Bugs, Cutworms, Aphids, Wheat fly.	40 - 50	15
Maize	Borers.	30 - 45	15

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours from last application.

PHYTOTOXICITY:

- Non-phytotoxic when used as recommended. Cotton, Brassicas, and some fruits may be injured by high rates of application.
- Russetting is possible for some apple varieties.

COMPATIBILITY:

Compatible with most insecticide except Benabacryl group and alkaline compounds, But compatibility test is recommended before 24 hours from spraying.

• For more details about first aid & precautions please refer to first aid & precautions index.