GARNET-5% SC

INSECTICIDE SUSPENSION CONCENTRATE



Contact and stomach broad spectrum insecticide, used to control many agricultural pests like fruit moth, vine moth, cabbage worm, beetles, thrips, leafminers and other insects on a wide range of agricultural crops listed in the table of uses.

COMPOSITION: Each litre contains:

FIPRONIL 5% W/V (A.I)

USES AND APPLICATION RATE:

| CROP | PESTS | APPLICATION RATE ML/200 LT | SAFETY PERIOD (DAYS) |
|---|--|--|----------------------------|
| Raddish, Cabbage, Lettuce, Cauliflower | Cabbage Worm. Diamond Back Moth. Cabbage Leaf Roller. | 50 (Foliar application) 25- 100 (Foliar application) 50 - 100 (Foliar application) | 7 |
| Eggplant | Thrips | 50 - 75 (Foliar application) | 14 |
| Grapevine | Vine Moth, Grape Berry Moth, Leafhopper | 100 - 200 (Foliar application) | 14 |
| Pepper | Pepper Weevil | 50 ml/ha | 14 |
| Potato | Wireworm. Potato Leafhopper. Thrips. | 200 (soil application). 200 (Foliar application) 50 - 100 (Foliar application) | |
| Peachs, Apple, Pear | Oriental Fruit Moth Plum Weevil. Codling Moth, Leafhopper. Leafminer, Psylla. | 100 ((Foliar application) 100 -200 (Foliar application) 100- 400 (Foliar application) 100 -300 (Foliar application) | 14 |

CAUTION:

Used only on above mentioned crops and follow the recommended safety period before harvest.

RE-ENTRY PERIOD:

24 hours from last application.

COMPATIBILITY:

Compatible with most pesticide, but compatibility test is recommended before spraying.

[•] For more details about first aid & precautions please refer to first aid & precautions index.