

# RATRYL-0.75% TP

RODENTICIDE  
READY TO USE POWDER

VAPCO

Class:  
Coumarin  
anticoagulant

Is an anticoagulant rodenticide which has shown exceptional activity against all rodent species, and can be used in all places where rodents can live.

**COMPOSITION:** Each kg contains:

**COUMATETRALYL**

**0.75% W/W (A.I)**

## PROPERTIES:

**RATRYL-0.75% TP:** It eliminates all type of mice and rats by eating small amounts over several consecutive days, as eating it only once does not give the desired results.

**RATRYL-0.75% TP:** It is acceptable for rodents that like its taste and does not cause them any caution when eating it.

**RATRYL-0.75% TP:** It causes rats a painless death like the natural death caused by old age without any other symptoms.

**RATRYL-0.75% TP:** It does not cause secondary poisoning to cats.

## USES AND RECOMMENDATIONS:

- Is recommended to control rats in cellars, farms, in stables housing large and small animals, in zoological gardens, slaughter-houses, cattle yards, goods depots and terminals, refuse tips, sewers, sewage works, port installations, on ships, in coal mines....etc.
- Is highly recommended for use in area-wide rat eradication programmes in cities, towns and rural districts. Control operations must be preceded by surveys to determine rat infestation density, and followed up to regular long-term inspections and re-treat to maintain the achieved state of clearance.
- Product may be used in control rats in agricultural crops.

## METHOD OF APPLICATION:

May be used with two methods:

- **First method: RATRYL-0.75% TP** is ready for use product used to control rats. It is applied by sprinkling it in a layer, in hiding places and on rat runways. The added amount must be regularly checked and replenished if necessary.
- **Second method: RATRYL-0.75% TP** can be mixed with cereals (Corn, Wheat, Barley ... etc) can be used by mixing 1 part of **RATRYL-0.75% TP** with 19 parts of bait material. The following mixture show exceptional effectiveness and it composed: 550 gm of ground Maize + 350 gm rolled Oats + 50 gm of castor Sugar + 50 gm of **RATRYL-0.75% TP**.

• For more details about first aid & precautions please refer to first aid & precautions index.

**REMARKS:**

- Do not touch **RATRYL-0.75% TP** with bare hands as human scent is a warning to rodents.
- Sprinkle the powder around rat hole and runways until signs of rat and mouse activity decrease.
- Do not place powder in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.
- Powder should not be touched after 5-8 days should be moved to a new location.
- Maintain an uninterrupted supply of fresh powder for 15 days or until all sign of feeding touching has disappeared.
- Remove all dead rats by placing them in tight-closing bags, after dusted with insecticide. The carceases should be handled with rubber or polyethylene gloves and all of the dead rats either burned or buried.
- Do not graze livestock in treated areas.
- Do not place the bait in open places.
- Do not place the baits in places easy to reach by children, livestock or pets and others.

• For more details about first aid & precautions please refer to first aid & precautions index.