



Fiprovap

(Pour-On)

COMPOSITION: each ml contains:

Fipronil 10 mg

INDICATIONS:

Treatment and control of the following external parasites of cattle, sheep, camelids, horses, guinea pigs: Ticks (*Boophilus microplus*), horn fly (*Haematobia irritans*), chewing lice (*Damalinia bovis*), sucking lice (*Linognathus vituli*) and nodular myiasis (*Dermatobia hominis*); flies (*Melophagus ovinus*).

DOSAGE AND METHOD OF APPLICATION

Epicutaneous application (pour-on):

- Cattle, sheep, camelids and horses: The recommended dose is 1 mL/10 kg of body weight, equivalent to 1 mg of fipronil/kg b.w.
- Guinea pigs: The recommended dose is of 0.1 mL/kg b.w., equivalent to 1 mg of fipronil/kg b.w.; which in the practice is equivalent to two drops per each kg b.w., applied on the back of the animal.

The treatment interval could be at least every two months, it will depend on the severity of the infestation, weather and management conditions in the herd. Consult your veterinarian to determine the better moment for the treatment.

WITHDRAWAL PERIOD

- Meat: 100 days.
- Do not use in lactating cattle producing milk for human consumption.
- Do not use in pregnant cows or heifers, which are intended to produce milk for human consumption, within 3 months of expected parturition.

PRECAUTIONS:

- External use only.
- Do not apply by another route of administration.
- Do not apply the product to wounded skin.
- Apply the product to the animal in a ventilated area.
- Do not use in sick or convalescent animals.

STORAGE:

- Keep away from Children, Food and drinks.
- Keep in dry, dark place below 30°C.

For Vet. Use only.