



Phoxivap

Emulsifiable Concentrate

COMPOSITION: Each ml contains:

Phoxime 500 mg

PROPERTIES:

Phoxime is an inhibitor of the enzyme cholinesterase at the synapses of the nerve ganglion. Inhibition of the enzyme is irreversible under physiological conditions.

INDICATIONS:

Treatment of myiasis, scabies and infestations by ticks, lice and keds in cattle, sheep, goats, horses.

DOSAGE:

By **bathing, spraying or rubbing with a sponge**, in a single application after dilution of the product according to the recommendations below:

	Sheep- Goat	Cattle	Equine
scabies			
Dose	1 lt in 2000 lt water. Old scabies: 1 lt in 1000 lt water.	1 lt in 1000 lt water. Old scabies 1 lt in 500 lt water.	1 ml in 1 lt water. Sponge application.
Keds, Lice, myiasis and Ticks.			
Dose	1 lt in 2000 lt water.	1 lt in 1000 lt water.	1 ml in 1 lt water Sponge application.

- Replenish: when the volume of the bath has decreased by 20%, refill with a solution of 250 ml of pure product in 125 liters water. Refill the bath the first 3 days and renew the complete bath on the 5th day.

WITHDRAWAL PERIOD:

Cattle:

- Meat and offal: 40 days.



- Milk: do not use in females producing milk for human consumption, lactating or in dry period.

Sheep, goats and horses:

- Meat and offal: 28 days.

Milk: do not use in females producing milk for human consumption, lactating or in dry period.

PRECAUTIONS:

- Do not use the drug 10 days before and 10 days after administration of other cholinesterase inhibitors, phenothiazines or muscle relaxants. Also avoid simultaneous use with general anesthetics.

STORAGE:

Keep in dry, dark place below 3°C.

For VET. Use only.