



VAZURIL 5%

(Suspension)

For Calves and Lambs

COMPOSITION: Each 1 ml contains:

Toltrazuril 50 mg

Indication:

- Lambs: The prevention of clinical signs of coccidiosis and reduction of coccidian shedding in Lambs on farms with a confirmed history of coccidiosis caused by *Eimeria crandallis* and *Eimeria ovinoidalis*.
- Calves: For the prevention of clinical signs of coccidiosis and reduction of coccidia shedding in housed calves replacing cows producing milk for human consumption (dairy cows) on farms with a confirmed history of coccidiosis caused by *Eimeria bovis* or *Eimeria zuernii*.

Dosage: Shake well before use.

- Lambs: Single oral dose of 0.4 ml oral suspension per kg body weight (20 mg toltrazuril/kg body weight).
- Calves: Single oral dose 3.0 ml oral suspension per 10 kg body weight (corresponding to 15 mg toltrazuril/kg body weight).

Withdrawal period:

- Calves: 63 days, not permitted for use in lactating animals producing milk for human consumption.
- Lambs: Meat and offal: 42 days, not permitted for use in lactating sheep producing milk for human consumption.

Contraindications:

- **Do not use in calves weighing more than 80 kg bodyweight.**
- **Do not use in fattening units such as veal or beef calves.**

Precautions:

- Frequent and repeated use of antiprotozoal from the same class may lead to the development of resistance.
- Avoid contact with skin, and dispose the unused drug.
- **Storage:**

Keep in dry, dark place below 30°C.

For vet use only.